

表 3371 透析前補正カルシウム濃度(mg/dL) 性別（透析患者全体）

性別	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
男性	733	1,144	4,478	14,573	4,156	45,814	51,973	32,419	3,857	9,697	4,615	1,488	516	359	175,822	22,319	198,141	9.14	0.73
(%)	(0.4)	(0.7)	(2.5)	(8.3)	(2.4)	(26.1)	(29.6)	(18.4)	(2.2)	(5.5)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
女性	409	505	1,921	5,866	1,716	21,505	30,031	21,550	2,741	7,280	3,655	1,293	457	322	99,251	12,716	111,967	9.28	0.76
(%)	(0.4)	(0.5)	(1.9)	(5.9)	(1.7)	(21.7)	(30.3)	(21.7)	(2.8)	(7.3)	(3.7)	(1.3)	(0.5)	(0.3)	(100.0)				
合計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし																			
(%)																			
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査