

表 3373 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別 (透析患者全体)

透析歴	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
2年未満	340	520	2,026	6,363	1,806	18,232	17,081	8,269	880	2,045	966	377	135	126	59,166	8,976	68,142	8.99	0.71
(%)	(0.6)	(0.9)	(3.4)	(10.8)	(3.1)	(30.8)	(28.9)	(14.0)	(1.5)	(3.5)	(1.6)	(0.6)	(0.2)	(0.2)	(100.0)				
2年～	239	358	1,469	5,159	1,513	18,216	21,705	12,922	1,436	3,474	1,556	523	165	145	68,880	9,063	77,943	9.14	0.70
(%)	(0.3)	(0.5)	(2.1)	(7.5)	(2.2)	(26.4)	(31.5)	(18.8)	(2.1)	(5.0)	(2.3)	(0.8)	(0.2)	(0.2)	(100.0)				
5年～	245	324	1,203	4,218	1,193	15,548	21,528	15,173	1,862	4,816	2,287	714	263	159	69,533	8,329	77,862	9.26	0.73
(%)	(0.4)	(0.5)	(1.7)	(6.1)	(1.7)	(22.4)	(31.0)	(21.8)	(2.7)	(6.9)	(3.3)	(1.0)	(0.4)	(0.2)	(100.0)				
10年～	126	144	623	2,033	574	7,151	10,250	8,469	1,110	3,016	1,578	527	208	122	35,931	4,101	40,032	9.34	0.77
(%)	(0.4)	(0.4)	(1.7)	(5.7)	(1.6)	(19.9)	(28.5)	(23.6)	(3.1)	(8.4)	(4.4)	(1.5)	(0.6)	(0.3)	(100.0)				
15年～	65	120	411	1,135	332	3,785	5,228	4,297	637	1,731	887	287	92	50	19,057	2,156	21,213	9.33	0.78
(%)	(0.3)	(0.6)	(2.2)	(6.0)	(1.7)	(19.9)	(27.4)	(22.5)	(3.3)	(9.1)	(4.7)	(1.5)	(0.5)	(0.3)	(100.0)				
20年～	57	75	310	716	215	2,090	2,969	2,313	324	898	479	148	57	33	10,684	1,118	11,802	9.28	0.81
(%)	(0.5)	(0.7)	(2.9)	(6.7)	(2.0)	(19.6)	(27.8)	(21.6)	(3.0)	(8.4)	(4.5)	(1.4)	(0.5)	(0.3)	(100.0)				
25年～	35	67	198	396	114	1,147	1,658	1,267	183	474	258	91	26	21	5,935	616	6,551	9.26	0.82
(%)	(0.6)	(1.1)	(3.3)	(6.7)	(1.9)	(19.3)	(27.9)	(21.3)	(3.1)	(8.0)	(4.3)	(1.5)	(0.4)	(0.4)	(100.0)				
30年～	21	24	112	250	69	691	981	768	97	319	155	71	13	14	3,585	375	3,960	9.28	0.82
(%)	(0.6)	(0.7)	(3.1)	(7.0)	(1.9)	(19.3)	(27.4)	(21.4)	(2.7)	(8.9)	(4.3)	(2.0)	(0.4)	(0.4)	(100.0)				
35年～	10	12	35	138	45	377	472	405	55	161	85	36	13	11	1,855	211	2,066	9.32	0.84
(%)	(0.5)	(0.6)	(1.9)	(7.4)	(2.4)	(20.3)	(25.4)	(21.8)	(3.0)	(8.7)	(4.6)	(1.9)	(0.7)	(0.6)	(100.0)				
40年～	3	5	10	29	11	67	117	78	12	36	17	7	1		393	58	451	9.26	0.79
(%)	(0.8)	(1.3)	(2.5)	(7.4)	(2.8)	(17.0)	(29.8)	(19.8)	(3.1)	(9.2)	(4.3)	(1.8)	(0.3)		(100.0)				
合計	1,141	1,649	6,397	20,437	5,872	67,304	81,989	53,961	6,596	16,970	8,268	2,781	973	681	275,019	35,003	310,022	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
不明	1		2			15	15	8	2	7	2				54	32	86	9.18	0.76
(%)	(1.9)		(3.7)			(27.8)	(27.8)	(14.8)	(3.7)	(13.0)	(3.7)				(100.0)				
記載なし																			
(%)																			
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	7.31	7.24	6.96	6.30	6.39	6.34	7.14	8.22	8.92	9.36	9.73	9.79	9.54	8.73	7.37	6.67	7.29		
標準偏差	8.39	8.54	8.19	7.50	7.59	7.10	7.29	7.54	7.76	7.93	8.00	8.43	7.94	8.38	7.52	7.26	7.49		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査