

表 3374 透析前補正カルシウム濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	362	492	1,897	5,696	1,620	18,971	25,089	18,435	2,395	6,393	3,092	1,052	356	234	86,084	10,886	96,970	9.25	0.76
(%)	(0.4)	(0.6)	(2.2)	(6.6)	(1.9)	(22.0)	(29.1)	(21.4)	(2.8)	(7.4)	(3.6)	(1.2)	(0.4)	(0.3)	(100.0)				
慢性腎盂腎炎	9	23	76	163	53	597	806	582	81	186	106	34	10	9	2,735	307	3,042	9.25	0.77
(%)	(0.3)	(0.8)	(2.8)	(6.0)	(1.9)	(21.8)	(29.5)	(21.3)	(3.0)	(6.8)	(3.9)	(1.2)	(0.4)	(0.3)	(100.0)				
急速進行性糸球体腎炎	10	9	33	138	37	568	702	445	51	118	56	13	6	2	2,188	306	2,494	9.18	0.69
(%)	(0.5)	(0.4)	(1.5)	(6.3)	(1.7)	(26.0)	(32.1)	(20.3)	(2.3)	(5.4)	(2.6)	(0.6)	(0.3)	(0.1)	(100.0)				
妊娠腎/妊娠中毒症	4	9	36	99	23	314	388	340	47	113	67	19	7	2	1,468	146	1,614	9.29	0.75
(%)	(0.3)	(0.6)	(2.5)	(6.7)	(1.6)	(21.4)	(26.4)	(23.2)	(3.2)	(7.7)	(4.6)	(1.3)	(0.5)	(0.1)	(100.0)				
その他の分類不能の腎炎	7	9	34	77	36	280	365	259	27	77	40	20	3	2	1,236	137	1,373	9.20	0.77
(%)	(0.6)	(0.7)	(2.8)	(6.2)	(2.9)	(22.7)	(29.5)	(21.0)	(2.2)	(6.2)	(3.2)	(1.6)	(0.2)	(0.2)	(100.0)				
多発性嚢胞腎	37	44	186	618	195	2,218	3,022	2,126	270	684	306	102	28	26	9,862	1,144	11,006	9.24	0.73
(%)	(0.4)	(0.4)	(1.9)	(6.3)	(2.0)	(22.5)	(30.6)	(21.6)	(2.7)	(6.9)	(3.1)	(1.0)	(0.3)	(0.3)	(100.0)				
腎硬化症	105	157	576	1,934	556	6,481	7,839	4,694	575	1,369	641	232	84	58	25,301	2,997	28,298	9.16	0.73
(%)	(0.4)	(0.6)	(2.3)	(7.6)	(2.2)	(25.6)	(31.0)	(18.6)	(2.3)	(5.4)	(2.5)	(0.9)	(0.3)	(0.2)	(100.0)				
悪性高血圧	9	18	68	154	68	541	691	439	46	142	58	24	9	2	2,269	304	2,573	9.16	0.73
(%)	(0.4)	(0.8)	(3.0)	(6.8)	(3.0)	(23.8)	(30.5)	(19.3)	(2.0)	(6.3)	(2.6)	(1.1)	(0.4)	(0.1)	(100.0)				
糖尿病性腎症	432	644	2,611	8,779	2,509	27,944	31,523	18,758	2,177	5,400	2,666	858	310	235	104,846	13,235	118,081	9.13	0.73
(%)	(0.4)	(0.6)	(2.5)	(8.4)	(2.4)	(26.7)	(30.1)	(17.9)	(2.1)	(5.2)	(2.5)	(0.8)	(0.3)	(0.2)	(100.0)				
SLE腎炎	3	15	37	138	31	459	576	410	60	141	71	20	10	4	1,975	261	2,236	9.25	0.75
(%)	(0.2)	(0.8)	(1.9)	(7.0)	(1.6)	(23.2)	(29.2)	(20.8)	(3.0)	(7.1)	(3.6)	(1.0)	(0.5)	(0.2)	(100.0)				
アミロイド腎	1	4	6	34	6	76	114	91	3	34	3	5	4	3	384	67	451	9.25	0.79
(%)	(0.3)	(1.0)	(1.6)	(8.9)	(1.6)	(19.8)	(29.7)	(23.7)	(0.8)	(8.9)	(0.8)	(1.3)	(1.0)	(0.8)	(100.0)				
痛風腎	8	4	19	68	20	215	279	235	22	83	32	8	1	3	997	116	1,113	9.24	0.76
(%)	(0.8)	(0.4)	(1.9)	(6.8)	(2.0)	(21.6)	(28.0)	(23.6)	(2.2)	(8.3)	(3.2)	(0.8)	(0.1)	(0.3)	(100.0)				
先天性代謝異常による腎不全	4	1	3	14	8	36	64	57	9	21	14	2	2		235	37	272	9.34	0.81
(%)	(1.7)	(0.4)	(1.3)	(6.0)	(3.4)	(15.3)	(27.2)	(24.3)	(3.8)	(8.9)	(6.0)	(0.9)	(0.9)		(100.0)				
腎・尿路結核	2		5	16	2	43	50	33	2	15	6	1	1		176	30	206	9.18	0.74
(%)	(1.1)		(2.8)	(9.1)	(1.1)	(24.4)	(28.4)	(18.8)	(1.1)	(8.5)	(3.4)	(0.6)	(0.6)		(100.0)				
腎・尿路結石	3	3	7	35	7	118	163	103	17	29	19	8	6	3	521	60	581	9.28	0.79
(%)	(0.6)	(0.6)	(1.3)	(6.7)	(1.3)	(22.6)	(31.3)	(19.8)	(3.3)	(5.6)	(3.6)	(1.5)	(1.2)	(0.6)	(100.0)				
腎・尿路腫瘍	3	5	14	64	13	205	232	142	20	38	32	8	2	3	781	105	886	9.18	0.75
(%)	(0.4)	(0.6)	(1.8)	(8.2)	(1.7)	(26.2)	(29.7)	(18.2)	(2.6)	(4.9)	(4.1)	(1.0)	(0.3)	(0.4)	(100.0)				
閉塞性尿路障害	4	3	15	47	16	159	167	135	24	39	14	7	2	3	635	88	723	9.19	0.76
(%)	(0.6)	(0.5)	(2.4)	(7.4)	(2.5)	(25.0)	(26.3)	(21.3)	(3.8)	(6.1)	(2.2)	(1.1)	(0.3)	(0.5)	(100.0)				
骨髄腫	4	5	1	24	4	64	56	42	6	14	9	1	1	1	232	44	276	9.10	0.86
(%)	(1.7)	(2.2)	(0.4)	(10.3)	(1.7)	(27.6)	(24.1)	(18.1)	(2.6)	(6.0)	(3.9)	(0.4)	(0.4)	(0.4)	(100.0)				
腎形成不全	4	7	19	36	13	128	145	128	18	38	22	7	5	4	574	75	649	9.24	0.86
(%)	(0.7)	(1.2)	(3.3)	(6.3)	(2.3)	(22.3)	(25.3)	(22.3)	(3.1)	(6.6)	(3.8)	(1.2)	(0.9)	(0.7)	(100.0)				
不明	88	145	556	1,717	485	5,874	7,162	4,769	543	1,480	753	277	95	63	24,007	3,537	27,544	9.20	0.75
(%)	(0.4)	(0.6)	(2.3)	(7.2)	(2.0)	(24.5)	(29.8)	(19.9)	(2.3)	(6.2)	(3.1)	(1.2)	(0.4)	(0.3)	(100.0)				
再導入	9	21	62	148	39	429	546	403	45	147	61	19	6	4	1,939	224	2,163	9.19	0.78
(%)	(0.5)	(1.1)	(3.2)	(7.6)	(2.0)	(22.1)	(28.2)	(20.8)	(2.3)	(7.6)	(3.1)	(1.0)	(0.3)	(0.2)	(100.0)				
その他	34	31	138	440	131	1,599	2,025	1,343	160	416	202	64	25	20	6,628	929	7,557	9.21	0.75
(%)	(0.5)	(0.5)	(2.1)	(6.6)	(2.0)	(24.1)	(30.6)	(20.3)	(2.4)	(6.3)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				
合計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし																			
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査