

表 3377 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別 (透析患者全体)

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	54	50	162	501	141	2,129	2,985	2,320	320	977	551	273	96	100	10,659	227	10,886	9.40	0.86
(%)	(0.5)	(0.5)	(1.5)	(4.7)	(1.3)	(20.0)	(28.0)	(21.8)	(3.0)	(9.2)	(5.2)	(2.6)	(0.9)	(0.9)	(100.0)				
3.0～	44	65	240	782	259	3,144	4,122	2,744	342	826	462	171	67	52	13,320	262	13,582	9.25	0.75
(%)	(0.3)	(0.5)	(1.8)	(5.9)	(1.9)	(23.6)	(30.9)	(20.6)	(2.6)	(6.2)	(3.5)	(1.3)	(0.5)	(0.4)	(100.0)				
3.5～	60	78	410	1,460	459	5,756	7,357	4,661	568	1,409	691	231	88	61	23,289	393	23,682	9.22	0.71
(%)	(0.3)	(0.3)	(1.8)	(6.3)	(2.0)	(24.7)	(31.6)	(20.0)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.3)	(100.0)				
4.0～	110	128	646	2,237	717	8,711	11,044	6,860	814	2,052	919	298	98	73	34,707	575	35,282	9.19	0.70
(%)	(0.3)	(0.4)	(1.9)	(6.4)	(2.1)	(25.1)	(31.8)	(19.8)	(2.3)	(5.9)	(2.6)	(0.9)	(0.3)	(0.2)	(100.0)				
4.5～	108	168	737	2,926	835	10,426	12,899	8,256	980	2,388	1,155	363	116	73	41,430	629	42,059	9.19	0.70
(%)	(0.3)	(0.4)	(1.8)	(7.1)	(2.0)	(25.2)	(31.1)	(19.9)	(2.4)	(5.8)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
5.0～	124	223	895	3,104	847	10,509	12,743	8,298	1,010	2,528	1,174	376	133	77	42,041	677	42,718	9.19	0.71
(%)	(0.3)	(0.5)	(2.1)	(7.4)	(2.0)	(25.0)	(30.3)	(19.7)	(2.4)	(6.0)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
5.5～	119	203	835	2,782	828	9,011	10,703	7,061	808	2,227	1,054	322	127	80	36,160	545	36,705	9.18	0.73
(%)	(0.3)	(0.6)	(2.3)	(7.7)	(2.3)	(24.9)	(29.6)	(19.5)	(2.2)	(6.2)	(2.9)	(0.9)	(0.4)	(0.2)	(100.0)				
6.0～	24	31	143	508	149	1,500	1,841	1,181	148	365	168	66	21	16	6,161	100	6,261	9.17	0.74
(%)	(0.4)	(0.5)	(2.3)	(8.2)	(2.4)	(24.3)	(29.9)	(19.2)	(2.4)	(5.9)	(2.7)	(1.1)	(0.3)	(0.3)	(100.0)				
6.1～	98	149	546	1,754	479	4,977	5,988	4,039	516	1,308	622	201	61	44	20,782	327	21,109	9.17	0.75
(%)	(0.5)	(0.7)	(2.6)	(8.4)	(2.3)	(23.9)	(28.8)	(19.4)	(2.5)	(6.3)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
6.5～	98	151	565	1,607	448	4,380	4,958	3,418	445	1,095	520	178	55	39	17,957	321	18,278	9.14	0.78
(%)	(0.5)	(0.8)	(3.1)	(8.9)	(2.5)	(24.4)	(27.6)	(19.0)	(2.5)	(6.1)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
7.0～	58	117	405	993	269	2,721	2,935	2,053	230	725	383	120	40	24	11,073	204	11,277	9.14	0.79
(%)	(0.5)	(1.1)	(3.7)	(9.0)	(2.4)	(24.6)	(26.5)	(18.5)	(2.1)	(6.5)	(3.5)	(1.1)	(0.4)	(0.2)	(100.0)				
7.5～	58	89	280	650	177	1,594	1,737	1,167	163	434	225	70	23	13	6,680	178	6,858	9.10	0.83
(%)	(0.9)	(1.3)	(4.2)	(9.7)	(2.6)	(23.9)	(26.0)	(17.5)	(2.4)	(6.5)	(3.4)	(1.0)	(0.3)	(0.2)	(100.0)				
8.0～	43	57	187	443	96	929	961	704	96	254	115	44	13	9	3,951	102	4,053	9.07	0.85
(%)	(1.1)	(1.4)	(4.7)	(11.2)	(2.4)	(23.5)	(24.3)	(17.8)	(2.4)	(6.4)	(2.9)	(1.1)	(0.3)	(0.2)	(100.0)				
8.5～	36	46	128	251	55	590	657	444	61	137	75	22	18	5	2,525	47	2,572	9.06	0.88
(%)	(1.4)	(1.8)	(5.1)	(9.9)	(2.2)	(23.4)	(26.0)	(17.6)	(2.4)	(5.4)	(3.0)	(0.9)	(0.7)	(0.2)	(100.0)				
9.0～	97	90	210	423	107	874	984	692	85	215	135	35	12	8	3,967	110	4,077	9.00	0.92
(%)	(2.4)	(2.3)	(5.3)	(10.7)	(2.7)	(22.0)	(24.8)	(17.4)	(2.1)	(5.4)	(3.4)	(0.9)	(0.3)	(0.2)	(100.0)				
合計	1,131	1,645	6,389	20,421	5,866	67,251	81,914	53,898	6,586	16,940	8,249	2,770	968	674	274,702	4,697	279,399	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし	11	4	10	18	6	68	90	71	12	37	21	11	5	7	371	30,338	30,709	9.37	1.16
(%)	(3.0)	(1.1)	(2.7)	(4.9)	(1.6)	(18.3)	(24.3)	(19.1)	(3.2)	(10.0)	(5.7)	(3.0)	(1.3)	(1.9)	(100.0)				
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	5.94	5.90	5.67	5.45	5.36	5.22	5.15	5.16	5.16	5.16	5.15	5.01	5.01	4.73	5.21	5.28	5.21		
標準偏差	2.05	1.76	1.64	1.47	1.42	1.38	1.37	1.40	1.44	1.47	1.54	1.60	1.62	1.75	1.42	1.58	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査