

表 3379 透析前補正カルシウム濃度(mg/dL) PTH値(pg/mL)別 (透析患者全体) PTH測定法:whole-PTH

PTH値	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満	7	14	32	93	26	241	490	461	61	203	130	50	13	21	1,842	52	1,894	9.50	0.86
(%)	(0.4)	(0.8)	(1.7)	(5.0)	(1.4)	(13.1)	(26.6)	(25.0)	(3.3)	(11.0)	(7.1)	(2.7)	(0.7)	(1.1)	(100.0)				
20～	11	4	19	99	37	429	766	565	60	162	77	24	9	6	2,268	63	2,331	9.33	0.70
(%)	(0.5)	(0.2)	(0.8)	(4.4)	(1.6)	(18.9)	(33.8)	(24.9)	(2.6)	(7.1)	(3.4)	(1.1)	(0.4)	(0.3)	(100.0)				
35～	3	7	18	104	29	423	696	455	48	110	46	9	6	4	1,958	85	2,043	9.25	0.63
(%)	(0.2)	(0.4)	(0.9)	(5.3)	(1.5)	(21.6)	(35.5)	(23.2)	(2.5)	(5.6)	(2.3)	(0.5)	(0.3)	(0.2)	(100.0)				
50～	5	5	35	165	50	752	1,094	642	60	156	76	15	3	3	3,061	92	3,153	9.21	0.62
(%)	(0.2)	(0.2)	(1.1)	(5.4)	(1.6)	(24.6)	(35.7)	(21.0)	(2.0)	(5.1)	(2.5)	(0.5)	(0.1)	(0.1)	(100.0)				
75～	7	5	38	194	60	712	876	511	46	143	46	11	6	1	2,656	92	2,748	9.15	0.64
(%)	(0.3)	(0.2)	(1.4)	(7.3)	(2.3)	(26.8)	(33.0)	(19.2)	(1.7)	(5.4)	(1.7)	(0.4)	(0.2)	(0.0)	(100.0)				
100～	8	13	40	201	60	612	681	392	52	115	38	10	1	1	2,224	74	2,298	9.09	0.66
(%)	(0.4)	(0.6)	(1.8)	(9.0)	(2.7)	(27.5)	(30.6)	(17.6)	(2.3)	(5.2)	(1.7)	(0.4)	(0.0)	(0.0)	(100.0)				
125～		10	39	122	39	424	447	277	21	79	28	8	1	1	1,496	43	1,539	9.11	0.66
(%)		(0.7)	(2.6)	(8.2)	(2.6)	(28.3)	(29.9)	(18.5)	(1.4)	(5.3)	(1.9)	(0.5)	(0.1)	(0.1)	(100.0)				
150～		3	1	10	18	10	18	10	1	3	1	1			48		48	9.27	0.66
(%)			(6.3)	(2.1)	(20.8)	(37.5)	(20.8)	(2.1)	(6.3)	(2.1)	(2.1)	(2.1)			(100.0)				
151～	8	8	27	115	36	320	360	242	23	63	26	5	1	3	1,237	26	1,263	9.10	0.72
(%)	(0.6)	(0.6)	(2.2)	(9.3)	(2.9)	(25.9)	(29.1)	(19.6)	(1.9)	(5.1)	(2.1)	(0.4)	(0.1)	(0.2)	(100.0)				
180～	3	7	32	90	19	213	260	161	18	32	18	2	1		856	19	875	9.06	0.69
(%)	(0.4)	(0.8)	(3.7)	(10.5)	(2.2)	(24.9)	(30.4)	(18.8)	(2.1)	(3.7)	(2.1)	(0.2)	(0.1)		(100.0)				
210～	5	7	22	62	11	154	172	110	18	29	13	5	3		611	17	628	9.09	0.75
(%)	(0.8)	(1.1)	(3.6)	(10.1)	(1.8)	(25.2)	(28.2)	(18.0)	(2.9)	(4.7)	(2.1)	(0.8)	(0.5)		(100.0)				
240～	10	4	12	70	22	166	190	121	17	40	11	4			667	7	674	9.07	0.73
(%)	(1.5)	(0.6)	(1.8)	(10.5)	(3.3)	(24.9)	(28.5)	(18.1)	(2.5)	(6.0)	(1.6)	(0.6)			(100.0)				
300～	1	4	7	31	8	73	67	49	12	16	6	4	4		282	2	284	9.13	0.79
(%)	(0.4)	(1.4)	(2.5)	(11.0)	(2.8)	(25.9)	(23.8)	(17.4)	(4.3)	(5.7)	(2.1)	(1.4)	(1.4)		(100.0)				
350～	2	1	10	20	4	35	39	37	4	15	7				174	5	179	9.11	0.81
(%)	(1.1)	(0.6)	(5.7)	(11.5)	(2.3)	(20.1)	(22.4)	(21.3)	(2.3)	(8.6)	(4.0)				(100.0)				
400～	3	2	10	18	5	47	44	47	2	12	4	5		1	200	2	202	9.11	0.90
(%)	(1.5)	(1.0)	(5.0)	(9.0)	(2.5)	(23.5)	(22.0)	(23.5)	(1.0)	(6.0)	(2.0)	(2.5)		(0.5)	(100.0)				
500～	4	1	12	20	3	59	60	46	10	18	8	1	2	1	245	2	247	9.16	0.84
(%)	(1.6)	(0.4)	(4.9)	(8.2)	(1.2)	(24.1)	(24.5)	(18.8)	(4.1)	(7.3)	(3.3)	(0.4)	(0.8)	(0.4)	(100.0)				
合計	77	92	356	1,405	409	4,670	6,260	4,126	453	1,196	535	154	50	42	19,825	581	20,406	9.20	0.70
(%)	(0.4)	(0.5)	(1.8)	(7.1)	(2.1)	(23.6)	(31.6)	(20.8)	(2.3)	(6.0)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
記載なし		1	5	20	10	69	96	66	12	19	9	6	4	2	319	338	657	9.29	0.76
(%)		(0.3)	(1.6)	(6.3)	(3.1)	(21.6)	(30.1)	(20.7)	(3.8)	(6.0)	(2.8)	(1.9)	(1.3)	(0.6)	(100.0)				
総計	77	93	361	1,425	419	4,739	6,356	4,192	465	1,215	544	160	54	44	20,144	919	21,063	9.20	0.70
(%)	(0.4)	(0.5)	(1.8)	(7.1)	(2.1)	(23.5)	(31.6)	(20.8)	(2.3)	(6.0)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
平均	180.09	136.21	152.46	131.23	118.63	116.06	101.33	101.82	111.65	101.14	91.25	86.53	107.76	66.98	108.54	89.65	108.01		
標準偏差	206.89	107.19	137.36	120.07	92.99	115.03	99.24	121.83	135.98	120.53	131.56	111.15	172.16	142.52	114.88	76.78	114.01		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査