

表 3380 透析前補正カルシウム濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別（透析患者全体）

透析前ヘモグロビン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
5未満	5		7	29	4	75	73	57	7	21	17	2	3		300	3	303	9.19	0.83
(%)	(1.7)		(2.3)	(9.7)	(1.3)	(25.0)	(24.3)	(19.0)	(2.3)	(7.0)	(5.7)	(0.7)	(1.0)		(100.0)				
5～	3	7	19	38	15	85	115	80	6	30	14	3	3	4	422	7	429	9.13	0.91
(%)	(0.7)	(1.7)	(4.5)	(9.0)	(3.6)	(20.1)	(27.3)	(19.0)	(1.4)	(7.1)	(3.3)	(0.7)	(0.7)	(0.9)	(100.0)				
6～	12	8	22	70	26	236	284	205	35	97	59	29	15	16	1,114	46	1,160	9.36	0.95
(%)	(1.1)	(0.7)	(2.0)	(6.3)	(2.3)	(21.2)	(25.5)	(18.4)	(3.1)	(8.7)	(5.3)	(2.6)	(1.3)	(1.4)	(100.0)				
7～	34	36	111	329	88	897	1,149	808	114	356	182	97	39	36	4,276	159	4,435	9.28	0.92
(%)	(0.8)	(0.8)	(2.6)	(7.7)	(2.1)	(21.0)	(26.9)	(18.9)	(2.7)	(8.3)	(4.3)	(2.3)	(0.9)	(0.8)	(100.0)				
8～	85	102	395	1,056	270	3,116	3,865	2,779	332	1,072	622	257	110	87	14,148	406	14,554	9.26	0.84
(%)	(0.6)	(0.7)	(2.8)	(7.5)	(1.9)	(22.0)	(27.3)	(19.6)	(2.3)	(7.6)	(4.4)	(1.8)	(0.8)	(0.6)	(100.0)				
9～	193	291	1,160	3,365	958	10,707	12,950	8,850	1,082	2,951	1,535	574	199	147	44,962	1,027	45,989	9.21	0.77
(%)	(0.4)	(0.6)	(2.6)	(7.5)	(2.1)	(23.8)	(28.8)	(19.7)	(2.4)	(6.6)	(3.4)	(1.3)	(0.4)	(0.3)	(100.0)				
10～	324	519	2,053	6,842	1,997	22,947	28,104	18,205	2,203	5,528	2,515	755	252	144	92,388	1,524	93,912	9.17	0.72
(%)	(0.4)	(0.6)	(2.2)	(7.4)	(2.2)	(24.8)	(30.4)	(19.7)	(2.4)	(6.0)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
11～	288	407	1,646	5,687	1,645	19,457	23,611	15,122	1,789	4,463	2,064	625	203	134	77,141	1,032	78,173	9.17	0.71
(%)	(0.4)	(0.5)	(2.1)	(7.4)	(2.1)	(25.2)	(30.6)	(19.6)	(2.3)	(5.8)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
12～	116	180	658	2,103	627	7,015	8,475	5,528	719	1,700	841	274	97	69	28,402	452	28,854	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.2)	(24.7)	(29.8)	(19.5)	(2.5)	(6.0)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
13～	42	54	191	568	134	1,691	2,058	1,391	184	459	254	92	30	28	7,176	132	7,308	9.20	0.80
(%)	(0.6)	(0.8)	(2.7)	(7.9)	(1.9)	(23.6)	(28.7)	(19.4)	(2.6)	(6.4)	(3.5)	(1.3)	(0.4)	(0.4)	(100.0)				
14～	26	22	75	187	66	565	729	516	78	185	91	44	17	11	2,612	61	2,673	9.21	0.85
(%)	(1.0)	(0.8)	(2.9)	(7.2)	(2.5)	(21.6)	(27.9)	(19.8)	(3.0)	(7.1)	(3.5)	(1.7)	(0.7)	(0.4)	(100.0)				
合計	1,128	1,626	6,337	20,274	5,830	66,791	81,413	53,541	6,549	16,862	8,194	2,752	968	676	272,941	4,849	277,790	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし	14	23	62	165	42	528	591	428	49	115	76	29	5	5	2,132	30,186	32,318	9.17	0.77
(%)	(0.7)	(1.1)	(2.9)	(7.7)	(2.0)	(24.8)	(27.7)	(20.1)	(2.3)	(5.4)	(3.6)	(1.4)	(0.2)	(0.2)	(100.0)				
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	10.64	10.66	10.67	10.71	10.73	10.74	10.74	10.71	10.72	10.64	10.59	10.50	10.39	10.28	10.71	10.47	10.71		
標準偏差	1.63	1.38	1.36	1.28	1.28	1.24	1.25	1.28	1.31	1.34	1.43	1.52	1.63	1.72	1.28	1.44	1.28		

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査