

表 3381 透析前補正カルシウム濃度(mg/dL) 総コレステロール濃度(mg/dL)別 (透析患者全体)

総コレステロール 濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
100未満	23	36	179	565	156	1,891	2,166	1,500	182	547	283	118	54	41	7,741	89	7,830	9.25	0.80
(%)	(0.3)	(0.5)	(2.3)	(7.3)	(2.0)	(24.4)	(28.0)	(19.4)	(2.4)	(7.1)	(3.7)	(1.5)	(0.7)	(0.5)	(100.0)				
100～	99	110	461	1,810	542	5,690	6,594	4,222	504	1,387	669	258	86	59	22,491	231	22,722	9.18	0.75
(%)	(0.4)	(0.5)	(2.0)	(8.0)	(2.4)	(25.3)	(29.3)	(18.8)	(2.2)	(6.2)	(3.0)	(1.1)	(0.4)	(0.3)	(100.0)				
120～	133	238	1,002	3,232	913	10,758	12,568	7,668	984	2,427	1,235	394	142	89	41,783	418	42,201	9.17	0.73
(%)	(0.3)	(0.6)	(2.4)	(7.7)	(2.2)	(25.7)	(30.1)	(18.4)	(2.4)	(5.8)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
140～	171	287	1,076	3,721	1,079	12,157	14,370	9,383	1,116	2,843	1,338	424	132	92	48,189	470	48,659	9.17	0.72
(%)	(0.4)	(0.6)	(2.2)	(7.7)	(2.2)	(25.2)	(29.8)	(19.5)	(2.3)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
160～	177	241	944	2,904	834	9,748	11,904	8,023	967	2,461	1,084	377	120	82	39,866	377	40,243	9.18	0.74
(%)	(0.4)	(0.6)	(2.4)	(7.3)	(2.1)	(24.5)	(29.9)	(20.1)	(2.4)	(6.2)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
180～	105	162	563	1,747	476	6,027	7,676	5,245	661	1,513	777	212	75	41	25,280	245	25,525	9.20	0.73
(%)	(0.4)	(0.6)	(2.2)	(6.9)	(1.9)	(23.8)	(30.4)	(20.7)	(2.6)	(6.0)	(3.1)	(0.8)	(0.3)	(0.2)	(100.0)				
200～	67	89	326	906	233	3,023	4,212	2,798	349	932	429	136	48	33	13,581	142	13,723	9.22	0.74
(%)	(0.5)	(0.7)	(2.4)	(6.7)	(1.7)	(22.3)	(31.0)	(20.6)	(2.6)	(6.9)	(3.2)	(1.0)	(0.4)	(0.2)	(100.0)				
220～	25	35	163	411	118	1,309	1,792	1,284	163	414	190	61	24	18	6,007	64	6,071	9.23	0.75
(%)	(0.4)	(0.6)	(2.7)	(6.8)	(2.0)	(21.8)	(29.8)	(21.4)	(2.7)	(6.9)	(3.2)	(1.0)	(0.4)	(0.3)	(100.0)				
240～	12	16	64	151	40	445	693	561	69	170	102	27	10	11	2,371	29	2,400	9.29	0.80
(%)	(0.5)	(0.7)	(2.7)	(6.4)	(1.7)	(18.8)	(29.2)	(23.7)	(2.9)	(7.2)	(4.3)	(1.1)	(0.4)	(0.5)	(100.0)				
260～	12	15	41	93	31	304	458	358	36	115	65	23	5	8	1,564	22	1,586	9.27	0.82
(%)	(0.8)	(1.0)	(2.6)	(5.9)	(2.0)	(19.4)	(29.3)	(22.9)	(2.3)	(7.4)	(4.2)	(1.5)	(0.3)	(0.5)	(100.0)				
合計	824	1,229	4,819	15,540	4,422	51,352	62,433	41,042	5,031	12,809	6,172	2,030	696	474	208,873	2,087	210,960	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.6)	(29.9)	(19.6)	(2.4)	(6.1)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
記載なし	318	420	1,580	4,899	1,450	15,967	19,571	12,927	1,567	4,168	2,098	751	277	207	66,200	32,948	99,148	9.20	0.76
(%)	(0.5)	(0.6)	(2.4)	(7.4)	(2.2)	(24.1)	(29.6)	(19.5)	(2.4)	(6.3)	(3.2)	(1.1)	(0.4)	(0.3)	(100.0)				
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	159.84	159.10	156.98	154.08	153.50	154.27	156.25	157.50	157.43	156.75	156.35	154.01	152.08	154.06	155.86	155.90	155.86		
標準偏差	38.72	35.76	36.00	34.81	34.65	34.50	35.47	36.47	36.32	37.38	38.21	39.17	39.55	43.68	35.72	37.34	35.74		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査