

表 3382 透析前補正カルシウム濃度(mg/dL) HDL-C濃度(mg/dL)別（透析患者全体）

HDL-C濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満	1	5	14	39	8	129	198	163	19	86	58	40	13	17	790	6	796	9.59	0.99
(%)	(0.1)	(0.6)	(1.8)	(4.9)	(1.0)	(16.3)	(25.1)	(20.6)	(2.4)	(10.9)	(7.3)	(5.1)	(1.6)	(2.2)	(100.0)				
20～	54	90	369	1,244	349	3,894	4,511	2,904	336	929	474	177	70	53	15,454	169	15,623	9.18	0.76
(%)	(0.3)	(0.6)	(2.4)	(8.0)	(2.3)	(25.2)	(29.2)	(18.8)	(2.2)	(6.0)	(3.1)	(1.1)	(0.5)	(0.3)	(100.0)				
30～	156	285	1,063	3,545	970	11,445	13,497	8,826	1,077	2,622	1,241	407	140	104	45,378	469	45,847	9.17	0.73
(%)	(0.3)	(0.6)	(2.3)	(7.8)	(2.1)	(25.2)	(29.7)	(19.4)	(2.4)	(5.8)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
40～	195	343	1,168	3,777	1,085	12,730	15,216	9,922	1,215	3,047	1,460	457	145	77	50,837	515	51,352	9.17	0.72
(%)	(0.4)	(0.7)	(2.3)	(7.4)	(2.1)	(25.0)	(29.9)	(19.5)	(2.4)	(6.0)	(2.9)	(0.9)	(0.3)	(0.2)	(100.0)				
50～	113	197	844	2,675	793	9,218	11,481	7,247	886	2,247	1,018	327	97	52	37,195	366	37,561	9.18	0.71
(%)	(0.3)	(0.5)	(2.3)	(7.2)	(2.1)	(24.8)	(30.9)	(19.5)	(2.4)	(6.0)	(2.7)	(0.9)	(0.3)	(0.1)	(100.0)				
60～	81	119	485	1,566	455	5,343	6,785	4,431	506	1,365	622	173	54	42	22,027	209	22,236	9.19	0.71
(%)	(0.4)	(0.5)	(2.2)	(7.1)	(2.1)	(24.3)	(30.8)	(20.1)	(2.3)	(6.2)	(2.8)	(0.8)	(0.2)	(0.2)	(100.0)				
70～	38	49	248	764	223	2,650	3,289	2,239	260	667	320	91	25	18	10,881	111	10,992	9.19	0.71
(%)	(0.3)	(0.5)	(2.3)	(7.0)	(2.0)	(24.4)	(30.2)	(20.6)	(2.4)	(6.1)	(2.9)	(0.8)	(0.2)	(0.2)	(100.0)				
80～	19	33	119	350	95	1,144	1,656	1,119	121	321	147	42	14	8	5,188	41	5,229	9.20	0.71
(%)	(0.4)	(0.6)	(2.3)	(6.7)	(1.8)	(22.1)	(31.9)	(21.6)	(2.3)	(6.2)	(2.8)	(0.8)	(0.3)	(0.2)	(100.0)				
90～	7	10	57	154	44	524	735	492	60	146	65	11	5	2	2,312	19	2,331	9.21	0.69
(%)	(0.3)	(0.4)	(2.5)	(6.7)	(1.9)	(22.7)	(31.8)	(21.3)	(2.6)	(6.3)	(2.8)	(0.5)	(0.2)	(0.1)	(100.0)				
100～	11	10	42	134	49	425	609	416	59	139	57	29	5	5	1,990	11	2,001	9.23	0.76
(%)	(0.6)	(0.5)	(2.1)	(6.7)	(2.5)	(21.4)	(30.6)	(20.9)	(3.0)	(7.0)	(2.9)	(1.5)	(0.3)	(0.3)	(100.0)				
合計	675	1,141	4,409	14,248	4,071	47,502	57,977	37,759	4,539	11,569	5,462	1,754	568	378	192,052	1,916	193,968	9.18	0.72
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.7)	(30.2)	(19.7)	(2.4)	(6.0)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
記載なし	467	508	1,990	6,191	1,801	19,817	24,027	16,210	2,059	5,408	2,808	1,027	405	303	83,021	33,119	116,140	9.20	0.79
(%)	(0.6)	(0.6)	(2.4)	(7.5)	(2.2)	(23.9)	(28.9)	(19.5)	(2.5)	(6.5)	(3.4)	(1.2)	(0.5)	(0.4)	(100.0)				
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	49.30	47.60	48.50	48.05	48.45	48.39	49.07	49.17	49.08	49.03	48.39	47.28	45.97	44.83	48.76	47.85	48.75		
標準偏差	18.21	16.09	16.63	16.43	16.76	16.32	16.76	16.92	17.02	17.18	16.85	17.85	16.60	19.02	16.71	15.79	16.71		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査