

表 3383 透析前補正カルシウム濃度(mg/dL) Non-HDL-C濃度(mg/dL)別 (透析患者全体)

Non-HDL-C濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
50未満	14	21	78	292	77	878	955	589	88	220	92	50	17	12	3,383	27	3,410	9.18	0.77
(%)	(0.4)	(0.6)	(2.3)	(8.6)	(2.3)	(26.0)	(28.2)	(17.4)	(2.6)	(6.5)	(2.7)	(1.5)	(0.5)	(0.4)	(100.0)				
50～	45	70	342	1,213	395	4,000	4,796	3,075	360	957	467	155	51	32	15,958	143	16,101	9.18	0.72
(%)	(0.3)	(0.4)	(2.1)	(7.6)	(2.5)	(25.1)	(30.1)	(19.3)	(2.3)	(6.0)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
70～	104	195	804	2,685	751	8,960	10,547	6,765	786	2,108	1,019	331	121	74	35,250	319	35,569	9.18	0.73
(%)	(0.3)	(0.6)	(2.3)	(7.6)	(2.1)	(25.4)	(29.9)	(19.2)	(2.2)	(6.0)	(2.9)	(0.9)	(0.3)	(0.2)	(100.0)				
90～	133	219	1,000	3,219	921	10,468	12,875	8,047	970	2,468	1,163	360	95	67	42,005	373	42,378	9.17	0.71
(%)	(0.3)	(0.5)	(2.4)	(7.7)	(2.2)	(24.9)	(30.7)	(19.2)	(2.3)	(5.9)	(2.8)	(0.9)	(0.2)	(0.2)	(100.0)				
110～	134	226	736	2,410	687	8,450	10,193	6,865	834	1,992	954	292	96	59	33,928	275	34,203	9.18	0.72
(%)	(0.4)	(0.7)	(2.2)	(7.1)	(2.0)	(24.9)	(30.0)	(20.2)	(2.5)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
130～	72	141	491	1,510	406	5,001	6,228	4,244	532	1,277	593	184	64	38	20,781	182	20,963	9.19	0.73
(%)	(0.3)	(0.7)	(2.4)	(7.3)	(2.0)	(24.1)	(30.0)	(20.4)	(2.6)	(6.1)	(2.9)	(0.9)	(0.3)	(0.2)	(100.0)				
150～	37	71	267	722	204	2,429	3,230	2,180	267	655	284	97	29	24	10,496	94	10,590	9.20	0.72
(%)	(0.4)	(0.7)	(2.5)	(6.9)	(1.9)	(23.1)	(30.8)	(20.8)	(2.5)	(6.2)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
170～	29	34	129	342	89	994	1,359	993	137	321	173	46	12	10	4,668	54	4,722	9.22	0.77
(%)	(0.6)	(0.7)	(2.8)	(7.3)	(1.9)	(21.3)	(29.1)	(21.3)	(2.9)	(6.9)	(3.7)	(1.0)	(0.3)	(0.2)	(100.0)				
190～	8	16	45	129	43	400	524	393	39	113	60	20	5	6	1,801	8	1,809	9.21	0.76
(%)	(0.4)	(0.9)	(2.5)	(7.2)	(2.4)	(22.2)	(29.1)	(21.8)	(2.2)	(6.3)	(3.3)	(1.1)	(0.3)	(0.3)	(100.0)				
210～	9	9	29	68	20	201	344	234	25	85	52	17	8	8	1,109	15	1,124	9.30	0.86
(%)	(0.8)	(0.8)	(2.6)	(6.1)	(1.8)	(18.1)	(31.0)	(21.1)	(2.3)	(7.7)	(4.7)	(1.5)	(0.7)	(0.7)	(100.0)				
合計	585	1,002	3,921	12,590	3,593	41,781	51,051	33,385	4,038	10,196	4,857	1,552	498	330	169,379	1,490	170,869	9.18	0.72
(%)	(0.3)	(0.6)	(2.3)	(7.4)	(2.1)	(24.7)	(30.1)	(19.7)	(2.4)	(6.0)	(2.9)	(0.9)	(0.3)	(0.2)	(100.0)				
記載なし	557	647	2,478	7,849	2,279	25,538	30,953	20,584	2,560	6,781	3,413	1,229	475	351	105,694	33,545	139,239	9.20	0.77
(%)	(0.5)	(0.6)	(2.3)	(7.4)	(2.2)	(24.2)	(29.3)	(19.5)	(2.4)	(6.4)	(3.2)	(1.2)	(0.4)	(0.3)	(100.0)				
総計	1,142	1,649	6,399	20,439	5,872	67,319	82,004	53,969	6,598	16,977	8,270	2,781	973	681	275,073	35,035	310,108	9.19	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.5)	(29.8)	(19.6)	(2.4)	(6.2)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	111.84	111.56	108.26	105.90	105.13	105.87	107.12	108.29	108.50	107.80	108.19	106.71	106.00	109.41	107.08	107.26	107.08		
標準偏差	37.64	34.75	34.48	33.34	33.16	32.69	33.38	34.00	33.96	34.62	35.51	36.77	36.02	40.91	33.60	34.08	33.61		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査