

表 3389 透析前補正カルシウム濃度(mg/dL) 透析歴(年)別 (施設血液透析患者 透析回数週3回)

透析歴	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
2年未満	265	424	1,625	5,268	1,505	15,026	13,891	6,506	689	1,593	740	274	91	93	47,990	2,725	50,715	8.98	0.70
(%)	(0.6)	(0.9)	(3.4)	(11.0)	(3.1)	(31.3)	(28.9)	(13.6)	(1.4)	(3.3)	(1.5)	(0.6)	(0.2)	(0.2)	(100.0)				
2年～	190	285	1,203	4,225	1,201	14,844	17,633	10,261	1,162	2,718	1,210	400	129	109	55,570	2,719	58,289	9.13	0.70
(%)	(0.3)	(0.5)	(2.2)	(7.6)	(2.2)	(26.7)	(31.7)	(18.5)	(2.1)	(4.9)	(2.2)	(0.7)	(0.2)	(0.2)	(100.0)				
5年～	188	254	972	3,410	985	12,678	17,575	12,305	1,519	3,864	1,863	567	213	131	56,524	2,703	59,227	9.26	0.72
(%)	(0.3)	(0.4)	(1.7)	(6.0)	(1.7)	(22.4)	(31.1)	(21.8)	(2.7)	(6.8)	(3.3)	(1.0)	(0.4)	(0.2)	(100.0)				
10年～	96	110	471	1,595	451	5,683	8,158	6,830	898	2,432	1,271	441	170	103	28,709	1,323	30,032	9.35	0.77
(%)	(0.3)	(0.4)	(1.6)	(5.6)	(1.6)	(19.8)	(28.4)	(23.8)	(3.1)	(8.5)	(4.4)	(1.5)	(0.6)	(0.4)	(100.0)				
15年～	50	78	296	848	249	2,851	3,957	3,295	498	1,364	689	228	77	46	14,526	723	15,249	9.34	0.78
(%)	(0.3)	(0.5)	(2.0)	(5.8)	(1.7)	(19.6)	(27.2)	(22.7)	(3.4)	(9.4)	(4.7)	(1.6)	(0.5)	(0.3)	(100.0)				
20年～	37	57	213	494	158	1,447	2,070	1,654	237	667	359	116	47	29	7,585	346	7,931	9.30	0.82
(%)	(0.5)	(0.8)	(2.8)	(6.5)	(2.1)	(19.1)	(27.3)	(21.8)	(3.1)	(8.8)	(4.7)	(1.5)	(0.6)	(0.4)	(100.0)				
25年～	24	38	112	238	60	747	1,073	847	118	327	174	63	21	12	3,854	190	4,044	9.29	0.82
(%)	(0.6)	(1.0)	(2.9)	(6.2)	(1.6)	(19.4)	(27.8)	(22.0)	(3.1)	(8.5)	(4.5)	(1.6)	(0.5)	(0.3)	(100.0)				
30年～	18	11	63	123	42	386	579	444	61	199	92	48	10	8	2,084	99	2,183	9.31	0.84
(%)	(0.9)	(0.5)	(3.0)	(5.9)	(2.0)	(18.5)	(27.8)	(21.3)	(2.9)	(9.5)	(4.4)	(2.3)	(0.5)	(0.4)	(100.0)				
35年～	3	7	14	73	21	192	229	215	34	89	48	22	10	8	965	68	1,033	9.37	0.86
(%)	(0.3)	(0.7)	(1.5)	(7.6)	(2.2)	(19.9)	(23.7)	(22.3)	(3.5)	(9.2)	(5.0)	(2.3)	(1.0)	(0.8)	(100.0)				
40年～	2	2	1	16	4	35	61	41	9	21	6	6	1		205	16	221	9.34	0.79
(%)	(1.0)	(1.0)	(0.5)	(7.8)	(2.0)	(17.1)	(29.8)	(20.0)	(4.4)	(10.2)	(2.9)	(2.9)	(0.5)		(100.0)				
合計	873	1,266	4,970	16,290	4,676	53,889	65,226	42,398	5,225	13,274	6,452	2,165	769	539	218,012	10,912	228,924	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
不明	1		1			13	10	6	2	3	1				37	10	47	9.14	0.73
(%)	(2.7)		(2.7)			(35.1)	(27.0)	(16.2)	(5.4)	(8.1)	(2.7)				(100.0)				
記載なし																			
(%)																			
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	6.95	6.50	6.27	5.78	5.89	5.94	6.75	7.91	8.62	9.10	9.42	9.71	9.83	8.77	6.99	6.76	6.98		
標準偏差	8.04	7.84	7.42	6.86	6.94	6.59	6.79	7.08	7.38	7.53	7.54	8.11	7.89	7.96	7.03	7.18	7.04		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査