

表 3392 透析前補正カルシウム濃度(mg/dL) 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満				5	3	18	19	24	2	7	3		1		82	2	84	9.33	0.69
(%)				(6.1)	(3.7)	(22.0)	(23.2)	(29.3)	(2.4)	(8.5)	(3.7)		(1.2)		(100.0)				
20～	3	3	13	48	14	152	218	176	28	61	32	14	6	2	770	32	802	9.34	0.79
(%)	(0.4)	(0.4)	(1.7)	(6.2)	(1.8)	(19.7)	(28.3)	(22.9)	(3.6)	(7.9)	(4.2)	(1.8)	(0.8)	(0.3)	(100.0)				
30～	79	88	354	1,103	361	4,098	5,342	3,875	535	1,409	756	329	103	107	18,539	578	19,117	9.30	0.80
(%)	(0.4)	(0.5)	(1.9)	(5.9)	(1.9)	(22.1)	(28.8)	(20.9)	(2.9)	(7.6)	(4.1)	(1.8)	(0.6)	(0.6)	(100.0)				
40～	226	291	1,134	3,798	1,106	13,715	17,518	11,769	1,441	3,979	1,971	733	272	204	58,157	1,652	59,809	9.24	0.76
(%)	(0.4)	(0.5)	(1.9)	(6.5)	(1.9)	(23.6)	(30.1)	(20.2)	(2.5)	(6.8)	(3.4)	(1.3)	(0.5)	(0.4)	(100.0)				
50～	260	369	1,543	5,419	1,561	17,698	21,216	13,324	1,666	4,165	1,983	642	231	131	70,208	1,901	72,109	9.17	0.73
(%)	(0.4)	(0.5)	(2.2)	(7.7)	(2.2)	(25.2)	(30.2)	(19.0)	(2.4)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
60～	164	273	1,075	3,511	993	11,073	12,767	8,214	976	2,333	1,073	291	100	63	42,906	1,176	44,082	9.14	0.71
(%)	(0.4)	(0.6)	(2.5)	(8.2)	(2.3)	(25.8)	(29.8)	(19.1)	(2.3)	(5.4)	(2.5)	(0.7)	(0.2)	(0.1)	(100.0)				
70～	86	120	480	1,430	374	4,380	5,069	3,128	360	796	372	85	34	20	16,734	450	17,184	9.10	0.70
(%)	(0.5)	(0.7)	(2.9)	(8.5)	(2.2)	(26.2)	(30.3)	(18.7)	(2.2)	(4.8)	(2.2)	(0.5)	(0.2)	(0.1)	(100.0)				
80～	28	68	195	527	159	1,567	1,747	1,076	115	278	133	33	12	4	5,942	139	6,081	9.08	0.72
(%)	(0.5)	(1.1)	(3.3)	(8.9)	(2.7)	(26.4)	(29.4)	(18.1)	(1.9)	(4.7)	(2.2)	(0.6)	(0.2)	(0.1)	(100.0)				
90～	10	21	76	213	48	582	618	372	35	108	37	5	3	3	2,131	64	2,195	9.05	0.70
(%)	(0.5)	(1.0)	(3.6)	(10.0)	(2.3)	(27.3)	(29.0)	(17.5)	(1.6)	(5.1)	(1.7)	(0.2)	(0.1)	(0.1)	(100.0)				
100～	6	10	36	69	19	195	217	114	26	27	15	1	1		736	19	755	9.00	0.72
(%)	(0.8)	(1.4)	(4.9)	(9.4)	(2.6)	(26.5)	(29.5)	(15.5)	(3.5)	(3.7)	(2.0)	(0.1)	(0.1)		(100.0)				
110～	3	11	13	36	3	68	74	57	6	12	5	2			290	9	299	8.97	0.80
(%)	(1.0)	(3.8)	(4.5)	(12.4)	(1.0)	(23.4)	(25.5)	(19.7)	(2.1)	(4.1)	(1.7)	(0.7)			(100.0)				
120～	2	2	7	21	8	40	55	30	7	5	6		1		184	5	189	9.01	0.79
(%)	(1.1)	(1.1)	(3.8)	(11.4)	(4.3)	(21.7)	(29.9)	(16.3)	(3.8)	(2.7)	(3.3)		(0.5)		(100.0)				
合計	867	1,256	4,926	16,180	4,649	53,586	64,860	42,159	5,197	13,180	6,386	2,135	764	534	216,679	6,027	222,706	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.5)	(2.4)	(6.1)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし	7	10	45	110	27	316	376	245	30	97	67	30	5	5	1,370	4,895	6,265	9.23	0.82
(%)	(0.5)	(0.7)	(3.3)	(8.0)	(2.0)	(23.1)	(27.4)	(17.9)	(2.2)	(7.1)	(4.9)	(2.2)	(0.4)	(0.4)	(100.0)				
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	56.36	58.55	57.64	57.01	56.42	56.06	55.49	54.94	54.54	53.74	53.27	50.94	51.10	49.34	55.46	54.91	55.45		
標準偏差	14.30	15.16	13.95	13.21	13.17	12.89	12.79	12.77	12.96	12.41	12.48	11.45	11.74	11.34	12.90	12.89	12.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査