

表 3393 透析前補正カルシウム濃度(mg/dL) 透析前リン濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前リン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
3未満	44	42	133	427	110	1,726	2,470	1,915	261	807	460	216	77	84	8,772	189	8,961	9.41	0.86
(%)	(0.5)	(0.5)	(1.5)	(4.9)	(1.3)	(19.7)	(28.2)	(21.8)	(3.0)	(9.2)	(5.2)	(2.5)	(0.9)	(1.0)	(100.0)				
3.0～	35	50	189	619	201	2,530	3,349	2,216	283	665	378	136	57	42	10,750	230	10,980	9.26	0.75
(%)	(0.3)	(0.5)	(1.8)	(5.8)	(1.9)	(23.5)	(31.2)	(20.6)	(2.6)	(6.2)	(3.5)	(1.3)	(0.5)	(0.4)	(100.0)				
3.5～	52	66	328	1,194	364	4,725	5,891	3,701	443	1,089	520	187	73	53	18,686	332	19,018	9.21	0.71
(%)	(0.3)	(0.4)	(1.8)	(6.4)	(1.9)	(25.3)	(31.5)	(19.8)	(2.4)	(5.8)	(2.8)	(1.0)	(0.4)	(0.3)	(100.0)				
4.0～	77	86	503	1,789	570	6,970	8,919	5,428	649	1,613	728	226	76	59	27,693	499	28,192	9.19	0.70
(%)	(0.3)	(0.3)	(1.8)	(6.5)	(2.1)	(25.2)	(32.2)	(19.6)	(2.3)	(5.8)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
4.5～	80	141	569	2,342	658	8,424	10,284	6,485	772	1,848	884	291	91	53	32,922	546	33,468	9.19	0.70
(%)	(0.2)	(0.4)	(1.7)	(7.1)	(2.0)	(25.6)	(31.2)	(19.7)	(2.3)	(5.6)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
5.0～	95	173	693	2,508	692	8,386	10,094	6,433	790	1,983	907	288	111	66	33,219	590	33,809	9.18	0.71
(%)	(0.3)	(0.5)	(2.1)	(7.5)	(2.1)	(25.2)	(30.4)	(19.4)	(2.4)	(6.0)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
5.5～	96	157	637	2,228	672	7,167	8,470	5,506	648	1,742	826	248	96	56	28,549	463	29,012	9.17	0.72
(%)	(0.3)	(0.5)	(2.2)	(7.8)	(2.4)	(25.1)	(29.7)	(19.3)	(2.3)	(6.1)	(2.9)	(0.9)	(0.3)	(0.2)	(100.0)				
6.0～	16	22	117	400	123	1,191	1,442	920	115	280	138	53	14	15	4,846	82	4,928	9.17	0.74
(%)	(0.3)	(0.5)	(2.4)	(8.3)	(2.5)	(24.6)	(29.8)	(19.0)	(2.4)	(5.8)	(2.8)	(1.1)	(0.3)	(0.3)	(100.0)				
6.1～	74	105	405	1,364	385	3,968	4,670	3,123	402	1,020	476	162	45	33	16,232	273	16,505	9.17	0.75
(%)	(0.5)	(0.6)	(2.5)	(8.4)	(2.4)	(24.4)	(28.8)	(19.2)	(2.5)	(6.3)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
6.5～	67	116	449	1,262	355	3,436	3,900	2,700	351	850	400	138	43	30	14,097	280	14,377	9.14	0.77
(%)	(0.5)	(0.8)	(3.2)	(9.0)	(2.5)	(24.4)	(27.7)	(19.2)	(2.5)	(6.0)	(2.8)	(1.0)	(0.3)	(0.2)	(100.0)				
7.0～	42	89	314	789	207	2,158	2,296	1,607	180	566	291	87	33	19	8,678	182	8,860	9.13	0.78
(%)	(0.5)	(1.0)	(3.6)	(9.1)	(2.4)	(24.9)	(26.5)	(18.5)	(2.1)	(6.5)	(3.4)	(1.0)	(0.4)	(0.2)	(100.0)				
7.5～	46	64	228	498	134	1,263	1,367	904	132	327	169	48	17	7	5,204	155	5,359	9.09	0.81
(%)	(0.9)	(1.2)	(4.4)	(9.6)	(2.6)	(24.3)	(26.3)	(17.4)	(2.5)	(6.3)	(3.2)	(0.9)	(0.3)	(0.1)	(100.0)				
8.0～	35	42	144	341	79	748	752	560	74	189	91	31	9	7	3,102	90	3,192	9.07	0.84
(%)	(1.1)	(1.4)	(4.6)	(11.0)	(2.5)	(24.1)	(24.2)	(18.1)	(2.4)	(6.1)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
8.5～	28	36	99	197	45	472	505	348	49	109	61	17	15	2	1,983	40	2,023	9.05	0.88
(%)	(1.4)	(1.8)	(5.0)	(9.9)	(2.3)	(23.8)	(25.5)	(17.5)	(2.5)	(5.5)	(3.1)	(0.9)	(0.8)	(0.1)	(100.0)				
9.0～	76	75	160	322	77	695	769	508	69	162	106	28	8	6	3,061	95	3,156	9.00	0.92
(%)	(2.5)	(2.5)	(5.2)	(10.5)	(2.5)	(22.7)	(25.1)	(16.6)	(2.3)	(5.3)	(3.5)	(0.9)	(0.3)	(0.2)	(100.0)				
合計	863	1,264	4,968	16,280	4,672	53,859	65,178	42,354	5,218	13,250	6,435	2,156	765	532	217,794	4,046	221,840	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし	11	2	3	10	4	43	58	50	9	27	18	9	4	7	255	6,876	7,131	9.45	1.28
(%)	(4.3)	(0.8)	(1.2)	(3.9)	(1.6)	(16.9)	(22.7)	(19.6)	(3.5)	(10.6)	(7.1)	(3.5)	(1.6)	(2.7)	(100.0)				
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	5.93	5.90	5.66	5.43	5.35	5.21	5.13	5.14	5.15	5.14	5.13	4.99	4.98	4.64	5.20	5.28	5.20		
標準偏差	2.07	1.80	1.65	1.47	1.40	1.38	1.37	1.40	1.45	1.48	1.55	1.61	1.60	1.73	1.42	1.58	1.43		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査