

表 3395 透析前補正カルシウム濃度(mg/dL) PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満	4	12	22	60	17	179	390	379	54	170	105	41	10	16	1,459	40	1,499	9.54	0.84
(%)	(0.3)	(0.8)	(1.5)	(4.1)	(1.2)	(12.3)	(26.7)	(26.0)	(3.7)	(11.7)	(7.2)	(2.8)	(0.7)	(1.1)	(100.0)				
20～	9	2	14	69	32	351	608	454	44	123	56	20	7	2	1,791	57	1,848	9.32	0.69
(%)	(0.5)	(0.1)	(0.8)	(3.9)	(1.8)	(19.6)	(33.9)	(25.3)	(2.5)	(6.9)	(3.1)	(1.1)	(0.4)	(0.1)	(100.0)				
35～	3	5	16	83	17	336	551	357	35	86	37	7	4	4	1,541	72	1,613	9.25	0.63
(%)	(0.2)	(0.3)	(1.0)	(5.4)	(1.1)	(21.8)	(35.8)	(23.2)	(2.3)	(5.6)	(2.4)	(0.5)	(0.3)	(0.3)	(100.0)				
50～	5	5	25	126	37	591	880	482	45	116	53	11	2	2	2,380	79	2,459	9.19	0.62
(%)	(0.2)	(0.2)	(1.1)	(5.3)	(1.6)	(24.8)	(37.0)	(20.3)	(1.9)	(4.9)	(2.2)	(0.5)	(0.1)	(0.1)	(100.0)				
75～	7	4	31	157	48	580	671	388	38	98	38	7	6	1	2,074	74	2,148	9.13	0.65
(%)	(0.3)	(0.2)	(1.5)	(7.6)	(2.3)	(28.0)	(32.4)	(18.7)	(1.8)	(4.7)	(1.8)	(0.3)	(0.3)	(0.0)	(100.0)				
100～	6	10	34	160	51	489	520	294	40	80	26	8	1	1	1,720	67	1,787	9.07	0.65
(%)	(0.3)	(0.6)	(2.0)	(9.3)	(3.0)	(28.4)	(30.2)	(17.1)	(2.3)	(4.7)	(1.5)	(0.5)	(0.1)	(0.1)	(100.0)				
125～		8	32	93	32	329	350	200	18	62	21	5	1	1	1,152	35	1,187	9.10	0.66
(%)		(0.7)	(2.8)	(8.1)	(2.8)	(28.6)	(30.4)	(17.4)	(1.6)	(5.4)	(1.8)	(0.4)	(0.1)	(0.1)	(100.0)				
150～		3				9	13	8	1	2	1				37		37	9.24	0.65
(%)			(8.1)			(24.3)	(35.1)	(21.6)	(2.7)	(5.4)	(2.7)				(100.0)				
151～	6	8	21	85	29	240	254	184	16	48	19	3	1	1	915	19	934	9.08	0.72
(%)	(0.7)	(0.9)	(2.3)	(9.3)	(3.2)	(26.2)	(27.8)	(20.1)	(1.7)	(5.2)	(2.1)	(0.3)	(0.1)	(0.1)	(100.0)				
180～	2	6	21	73	17	158	198	122	15	22	15	1	1		651	15	666	9.06	0.69
(%)	(0.3)	(0.9)	(3.2)	(11.2)	(2.6)	(24.3)	(30.4)	(18.7)	(2.3)	(3.4)	(2.3)	(0.2)	(0.2)		(100.0)				
210～	4	7	17	46	9	118	125	83	17	22	9	1	3		461	11	472	9.07	0.76
(%)	(0.9)	(1.5)	(3.7)	(10.0)	(2.0)	(25.6)	(27.1)	(18.0)	(3.7)	(4.8)	(2.0)	(0.2)	(0.7)		(100.0)				
240～	8	4	9	61	19	128	145	91	15	31	7	3			521	5	526	9.06	0.73
(%)	(1.5)	(0.8)	(1.7)	(11.7)	(3.6)	(24.6)	(27.8)	(17.5)	(2.9)	(6.0)	(1.3)	(0.6)			(100.0)				
300～	1	3	4	19	2	53	51	35	9	14	5	3	4		203	2	205	9.20	0.80
(%)	(0.5)	(1.5)	(2.0)	(9.4)	(1.0)	(26.1)	(25.1)	(17.2)	(4.4)	(6.9)	(2.5)	(1.5)	(2.0)		(100.0)				
350～	2	1	8	17	3	26	27	25	4	10	7				130	5	135	9.08	0.86
(%)	(1.5)	(0.8)	(6.2)	(13.1)	(2.3)	(20.0)	(20.8)	(19.2)	(3.1)	(7.7)	(5.4)				(100.0)				
400～	3	1	9	10	4	42	32	31	1	9	3	4		1	150	2	152	9.08	0.94
(%)	(2.0)	(0.7)	(6.0)	(6.7)	(2.7)	(28.0)	(21.3)	(20.7)	(0.7)	(6.0)	(2.0)	(2.7)		(0.7)	(100.0)				
500～	4	1	8	15	2	40	37	26	8	13	7		2	1	164	2	166	9.15	0.91
(%)	(2.4)	(0.6)	(4.9)	(9.1)	(1.2)	(24.4)	(22.6)	(15.9)	(4.9)	(7.9)	(4.3)		(1.2)	(0.6)	(100.0)				
合計	64	77	274	1,074	319	3,669	4,852	3,159	360	906	409	114	42	30	15,349	485	15,834	9.19	0.71
(%)	(0.4)	(0.5)	(1.8)	(7.0)	(2.1)	(23.9)	(31.6)	(20.6)	(2.3)	(5.9)	(2.7)	(0.7)	(0.3)	(0.2)	(100.0)				
記載なし		1	4	16	8	60	84	51	11	16	8	4	2	1	266	172	438	9.26	0.72
(%)		(0.4)	(1.5)	(6.0)	(3.0)	(22.6)	(31.6)	(19.2)	(4.1)	(6.0)	(3.0)	(1.5)	(0.8)	(0.4)	(100.0)				
総計	64	78	278	1,090	327	3,729	4,936	3,210	371	922	417	118	44	31	15,615	657	16,272	9.19	0.71
(%)	(0.4)	(0.5)	(1.8)	(7.0)	(2.1)	(23.9)	(31.6)	(20.6)	(2.4)	(5.9)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
平均	191.52	139.48	151.85	131.72	118.80	114.55	98.43	96.71	112.64	97.74	93.16	74.75	123.33	74.83	106.23	88.91	105.70		
標準偏差	221.40	104.76	138.57	118.54	89.54	113.66	95.38	106.34	133.07	109.27	142.30	101.60	183.90	165.17	110.03	78.78	109.24		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査