

表 3396 透析前補正カルシウム濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
5未満	4		5	26	4	65	64	46	5	17	17	2	3		258	3	261	9.21	0.84
(%)	(1.6)		(1.9)	(10.1)	(1.6)	(25.2)	(24.8)	(17.8)	(1.9)	(6.6)	(6.6)	(0.8)	(1.2)		(100.0)				
5～	2	5	17	31	13	77	99	65	6	27	9	2	1	3	357	5	362	9.10	0.88
(%)	(0.6)	(1.4)	(4.8)	(8.7)	(3.6)	(21.6)	(27.7)	(18.2)	(1.7)	(7.6)	(2.5)	(0.6)	(0.3)	(0.8)	(100.0)				
6～	12	8	16	54	20	204	231	164	24	78	47	23	10	13	904	38	942	9.33	0.95
(%)	(1.3)	(0.9)	(1.8)	(6.0)	(2.2)	(22.6)	(25.6)	(18.1)	(2.7)	(8.6)	(5.2)	(2.5)	(1.1)	(1.4)	(100.0)				
7～	23	26	92	276	73	727	928	633	94	297	136	79	34	33	3,451	127	3,578	9.29	0.92
(%)	(0.7)	(0.8)	(2.7)	(8.0)	(2.1)	(21.1)	(26.9)	(18.3)	(2.7)	(8.6)	(3.9)	(2.3)	(1.0)	(1.0)	(100.0)				
8～	67	81	325	862	214	2,508	3,103	2,254	268	855	494	205	87	74	11,397	329	11,726	9.26	0.84
(%)	(0.6)	(0.7)	(2.9)	(7.6)	(1.9)	(22.0)	(27.2)	(19.8)	(2.4)	(7.5)	(4.3)	(1.8)	(0.8)	(0.6)	(100.0)				
9～	163	226	910	2,680	762	8,701	10,428	7,148	871	2,349	1,198	449	164	109	36,158	880	37,038	9.20	0.77
(%)	(0.5)	(0.6)	(2.5)	(7.4)	(2.1)	(24.1)	(28.8)	(19.8)	(2.4)	(6.5)	(3.3)	(1.2)	(0.5)	(0.3)	(100.0)				
10～	248	411	1,585	5,488	1,603	18,462	22,649	14,455	1,755	4,343	1,974	595	194	116	73,878	1,339	75,217	9.17	0.71
(%)	(0.3)	(0.6)	(2.1)	(7.4)	(2.2)	(25.0)	(30.7)	(19.6)	(2.4)	(5.9)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
11～	213	306	1,275	4,499	1,305	15,464	18,574	11,746	1,399	3,464	1,623	473	159	101	60,601	878	61,479	9.17	0.71
(%)	(0.4)	(0.5)	(2.1)	(7.4)	(2.2)	(25.5)	(30.6)	(19.4)	(2.3)	(5.7)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
12～	83	129	507	1,648	495	5,517	6,601	4,189	566	1,278	641	210	77	53	21,994	376	22,370	9.18	0.73
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.3)	(25.1)	(30.0)	(19.0)	(2.6)	(5.8)	(2.9)	(1.0)	(0.4)	(0.2)	(100.0)				
13～	29	36	141	437	103	1,328	1,548	999	146	347	188	72	23	23	5,420	117	5,537	9.19	0.80
(%)	(0.5)	(0.7)	(2.6)	(8.1)	(1.9)	(24.5)	(28.6)	(18.4)	(2.7)	(6.4)	(3.5)	(1.3)	(0.4)	(0.4)	(100.0)				
14～	17	18	50	148	50	408	522	372	54	132	61	29	15	9	1,885	48	1,933	9.21	0.84
(%)	(0.9)	(1.0)	(2.7)	(7.9)	(2.7)	(21.6)	(27.7)	(19.7)	(2.9)	(7.0)	(3.2)	(1.5)	(0.8)	(0.5)	(100.0)				
合計	861	1,246	4,923	16,149	4,642	53,461	64,747	42,071	5,188	13,187	6,388	2,139	767	534	216,303	4,140	220,443	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.5)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし	13	20	48	141	34	441	489	333	39	90	65	26	2	5	1,746	6,782	8,528	9.16	0.78
(%)	(0.7)	(1.1)	(2.7)	(8.1)	(1.9)	(25.3)	(28.0)	(19.1)	(2.2)	(5.2)	(3.7)	(1.5)	(0.1)	(0.3)	(100.0)				
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	10.56	10.63	10.64	10.69	10.72	10.72	10.72	10.69	10.71	10.61	10.57	10.46	10.39	10.24	10.69	10.47	10.69		
標準偏差	1.61	1.37	1.35	1.29	1.28	1.24	1.24	1.27	1.30	1.34	1.41	1.51	1.63	1.75	1.27	1.41	1.28		

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査