

表 3397 透析前補正カルシウム濃度(mg/dL) 総コレステロール濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

総コレステロール濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
100未満	19	28	141	459	131	1,596	1,789	1,220	145	449	231	95	44	32	6,379	75	6,454	9.24	0.80
(%)	(0.3)	(0.4)	(2.2)	(7.2)	(2.1)	(25.0)	(28.0)	(19.1)	(2.3)	(7.0)	(3.6)	(1.5)	(0.7)	(0.5)	(100.0)				
100～	77	84	380	1,485	437	4,692	5,419	3,431	422	1,130	537	203	71	51	18,419	201	18,620	9.17	0.75
(%)	(0.4)	(0.5)	(2.1)	(8.1)	(2.4)	(25.5)	(29.4)	(18.6)	(2.3)	(6.1)	(2.9)	(1.1)	(0.4)	(0.3)	(100.0)				
120～	100	192	813	2,640	751	8,811	10,221	6,208	804	1,963	964	317	114	73	33,971	344	34,315	9.16	0.73
(%)	(0.3)	(0.6)	(2.4)	(7.8)	(2.2)	(25.9)	(30.1)	(18.3)	(2.4)	(5.8)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
140～	136	227	849	3,008	868	9,764	11,480	7,393	907	2,239	1,044	338	100	78	38,431	395	38,826	9.17	0.72
(%)	(0.4)	(0.6)	(2.2)	(7.8)	(2.3)	(25.4)	(29.9)	(19.2)	(2.4)	(5.8)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
160～	137	172	706	2,299	663	7,749	9,424	6,221	737	1,859	855	287	97	64	31,270	313	31,583	9.18	0.73
(%)	(0.4)	(0.6)	(2.3)	(7.4)	(2.1)	(24.8)	(30.1)	(19.9)	(2.4)	(5.9)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
180～	70	127	397	1,344	376	4,719	5,901	3,998	515	1,142	589	158	53	32	19,421	198	19,619	9.20	0.72
(%)	(0.4)	(0.7)	(2.0)	(6.9)	(1.9)	(24.3)	(30.4)	(20.6)	(2.7)	(5.9)	(3.0)	(0.8)	(0.3)	(0.2)	(100.0)				
200～	52	66	254	661	183	2,292	3,146	2,042	272	654	316	106	34	25	10,103	106	10,209	9.21	0.75
(%)	(0.5)	(0.7)	(2.5)	(6.5)	(1.8)	(22.7)	(31.1)	(20.2)	(2.7)	(6.5)	(3.1)	(1.0)	(0.3)	(0.2)	(100.0)				
220～	21	26	119	285	80	1,001	1,338	922	122	300	137	38	21	15	4,425	50	4,475	9.23	0.75
(%)	(0.5)	(0.6)	(2.7)	(6.4)	(1.8)	(22.6)	(30.2)	(20.8)	(2.8)	(6.8)	(3.1)	(0.9)	(0.5)	(0.3)	(100.0)				
240～	7	12	50	113	31	335	483	406	46	111	80	18	9	7	1,708	25	1,733	9.27	0.79
(%)	(0.4)	(0.7)	(2.9)	(6.6)	(1.8)	(19.6)	(28.3)	(23.8)	(2.7)	(6.5)	(4.7)	(1.1)	(0.5)	(0.4)	(100.0)				
260～	6	12	27	72	25	216	323	243	24	87	40	17	3	1	1,096	15	1,111	9.24	0.78
(%)	(0.5)	(1.1)	(2.5)	(6.6)	(2.3)	(19.7)	(29.5)	(22.2)	(2.2)	(7.9)	(3.6)	(1.6)	(0.3)	(0.1)	(100.0)				
合計	625	946	3,736	12,366	3,545	41,175	49,524	32,084	3,994	9,934	4,793	1,577	546	378	165,223	1,722	166,945	9.18	0.73
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.9)	(30.0)	(19.4)	(2.4)	(6.0)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
記載なし	249	320	1,235	3,924	1,131	12,727	15,712	10,320	1,233	3,343	1,660	588	223	161	52,826	9,200	62,026	9.20	0.76
(%)	(0.5)	(0.6)	(2.3)	(7.4)	(2.1)	(24.1)	(29.7)	(19.5)	(2.3)	(6.3)	(3.1)	(1.1)	(0.4)	(0.3)	(100.0)				
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	158.29	158.54	155.63	152.96	152.65	153.35	155.13	156.21	156.23	155.16	155.34	152.68	151.36	150.99	154.71	154.89	154.71		
標準偏差	37.73	35.72	35.71	34.19	34.39	34.20	35.00	35.99	35.81	37.07	37.60	38.56	39.61	39.49	35.27	36.85	35.28		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査