

表 3398 透析前補正カルシウム濃度(mg/dL) HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

HDL-C濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満	1	4	13	36	6	107	164	136	18	73	49	34	10	15	666	5	671	9.58	1.00
(%)	(0.2)	(0.6)	(2.0)	(5.4)	(0.9)	(16.1)	(24.6)	(20.4)	(2.7)	(11.0)	(7.4)	(5.1)	(1.5)	(2.3)	(100.0)				
20～	43	71	287	1,037	292	3,242	3,732	2,363	269	750	384	136	57	44	12,707	139	12,846	9.17	0.75
(%)	(0.3)	(0.6)	(2.3)	(8.2)	(2.3)	(25.5)	(29.4)	(18.6)	(2.1)	(5.9)	(3.0)	(1.1)	(0.4)	(0.3)	(100.0)				
30～	121	215	838	2,900	785	9,252	10,869	7,070	876	2,066	979	331	116	85	36,503	385	36,888	9.17	0.73
(%)	(0.3)	(0.6)	(2.3)	(7.9)	(2.2)	(25.3)	(29.8)	(19.4)	(2.4)	(5.7)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
40～	139	277	912	2,998	884	10,176	12,150	7,764	966	2,392	1,136	345	112	61	40,312	417	40,729	9.17	0.72
(%)	(0.3)	(0.7)	(2.3)	(7.4)	(2.2)	(25.2)	(30.1)	(19.3)	(2.4)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
50～	84	150	654	2,067	627	7,202	8,959	5,563	680	1,693	756	244	71	37	28,787	293	29,080	9.17	0.70
(%)	(0.3)	(0.5)	(2.3)	(7.2)	(2.2)	(25.0)	(31.1)	(19.3)	(2.4)	(5.9)	(2.6)	(0.8)	(0.2)	(0.1)	(100.0)				
60～	65	90	369	1,185	346	4,066	5,143	3,294	387	1,020	459	134	41	33	16,632	160	16,792	9.18	0.71
(%)	(0.4)	(0.5)	(2.2)	(7.1)	(2.1)	(24.4)	(30.9)	(19.8)	(2.3)	(6.1)	(2.8)	(0.8)	(0.2)	(0.2)	(100.0)				
70～	28	32	173	570	167	2,035	2,423	1,662	195	485	240	68	23	13	8,114	87	8,201	9.19	0.71
(%)	(0.3)	(0.4)	(2.1)	(7.0)	(2.1)	(25.1)	(29.9)	(20.5)	(2.4)	(6.0)	(3.0)	(0.8)	(0.3)	(0.2)	(100.0)				
80～	14	20	80	271	67	842	1,218	805	96	232	111	32	10	7	3,805	29	3,834	9.21	0.71
(%)	(0.4)	(0.5)	(2.1)	(7.1)	(1.8)	(22.1)	(32.0)	(21.2)	(2.5)	(6.1)	(2.9)	(0.8)	(0.3)	(0.2)	(100.0)				
90～	5	8	48	125	30	384	547	362	40	99	51	10	5	1	1,715	18	1,733	9.20	0.69
(%)	(0.3)	(0.5)	(2.8)	(7.3)	(1.7)	(22.4)	(31.9)	(21.1)	(2.3)	(5.8)	(3.0)	(0.6)	(0.3)	(0.1)	(100.0)				
100～	9	7	34	102	37	326	442	287	44	93	38	22	5	4	1,450	8	1,458	9.20	0.78
(%)	(0.6)	(0.5)	(2.3)	(7.0)	(2.6)	(22.5)	(30.5)	(19.8)	(3.0)	(6.4)	(2.6)	(1.5)	(0.3)	(0.3)	(100.0)				
合計	509	874	3,408	11,291	3,241	37,632	45,647	29,306	3,571	8,903	4,203	1,356	450	300	150,691	1,541	152,232	9.18	0.72
(%)	(0.3)	(0.6)	(2.3)	(7.5)	(2.2)	(25.0)	(30.3)	(19.4)	(2.4)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
記載なし	365	392	1,563	4,999	1,435	16,270	19,589	13,098	1,656	4,374	2,250	809	319	239	67,358	9,381	76,739	9.20	0.78
(%)	(0.5)	(0.6)	(2.3)	(7.4)	(2.1)	(24.2)	(29.1)	(19.4)	(2.5)	(6.5)	(3.3)	(1.2)	(0.5)	(0.4)	(100.0)				
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	49.35	47.05	48.16	47.61	47.92	47.95	48.56	48.62	48.59	48.40	47.90	47.06	45.96	44.36	48.26	47.54	48.25		
標準偏差	18.88	15.40	16.61	16.39	16.60	16.19	16.56	16.71	16.80	16.92	16.78	18.03	17.10	19.44	16.55	15.78	16.54		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査