

表 3399 透析前補正カルシウム濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

Non-HDL-C濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
50未満	12	13	60	238	61	720	754	463	68	177	67	38	14	8	2,693	23	2,716	9.17	0.77
(%)	(0.4)	(0.5)	(2.2)	(8.8)	(2.3)	(26.7)	(28.0)	(17.2)	(2.5)	(6.6)	(2.5)	(1.4)	(0.5)	(0.3)	(100.0)				
50～	36	60	268	954	311	3,243	3,846	2,464	294	750	371	122	43	29	12,791	115	12,906	9.18	0.72
(%)	(0.3)	(0.5)	(2.1)	(7.5)	(2.4)	(25.4)	(30.1)	(19.3)	(2.3)	(5.9)	(2.9)	(1.0)	(0.3)	(0.2)	(100.0)				
70～	80	156	670	2,184	617	7,223	8,476	5,332	638	1,668	795	266	94	64	28,263	269	28,532	9.17	0.73
(%)	(0.3)	(0.6)	(2.4)	(7.7)	(2.2)	(25.6)	(30.0)	(18.9)	(2.3)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
90～	102	166	758	2,585	743	8,371	10,212	6,313	769	1,946	892	282	78	56	33,273	306	33,579	9.17	0.71
(%)	(0.3)	(0.5)	(2.3)	(7.8)	(2.2)	(25.2)	(30.7)	(19.0)	(2.3)	(5.8)	(2.7)	(0.8)	(0.2)	(0.2)	(100.0)				
110～	110	164	564	1,918	537	6,690	7,966	5,293	651	1,475	748	225	79	43	26,463	216	26,679	9.17	0.72
(%)	(0.4)	(0.6)	(2.1)	(7.2)	(2.0)	(25.3)	(30.1)	(20.0)	(2.5)	(5.6)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
130～	47	109	369	1,169	333	3,914	4,833	3,184	401	971	439	135	51	32	15,987	146	16,133	9.18	0.72
(%)	(0.3)	(0.7)	(2.3)	(7.3)	(2.1)	(24.5)	(30.2)	(19.9)	(2.5)	(6.1)	(2.7)	(0.8)	(0.3)	(0.2)	(100.0)				
150～	29	63	202	552	152	1,882	2,465	1,624	199	471	215	73	22	17	7,966	77	8,043	9.19	0.72
(%)	(0.4)	(0.8)	(2.5)	(6.9)	(1.9)	(23.6)	(30.9)	(20.4)	(2.5)	(5.9)	(2.7)	(0.9)	(0.3)	(0.2)	(100.0)				
170～	22	31	100	249	69	763	1,039	734	108	226	137	30	8	8	3,524	43	3,567	9.21	0.76
(%)	(0.6)	(0.9)	(2.8)	(7.1)	(2.0)	(21.7)	(29.5)	(20.8)	(3.1)	(6.4)	(3.9)	(0.9)	(0.2)	(0.2)	(100.0)				
190～	5	11	35	100	32	287	381	292	34	90	44	13	3	4	1,331	7	1,338	9.22	0.76
(%)	(0.4)	(0.8)	(2.6)	(7.5)	(2.4)	(21.6)	(28.6)	(21.9)	(2.6)	(6.8)	(3.3)	(1.0)	(0.2)	(0.3)	(100.0)				
210～	4	5	19	52	17	152	264	176	16	59	33	16	7	2	822	10	832	9.29	0.79
(%)	(0.5)	(0.6)	(2.3)	(6.3)	(2.1)	(18.5)	(32.1)	(21.4)	(1.9)	(7.2)	(4.0)	(1.9)	(0.9)	(0.2)	(100.0)				
合計	447	778	3,045	10,001	2,872	33,245	40,236	25,875	3,178	7,833	3,741	1,200	399	263	133,113	1,212	134,325	9.18	0.72
(%)	(0.3)	(0.6)	(2.3)	(7.5)	(2.2)	(25.0)	(30.2)	(19.4)	(2.4)	(5.9)	(2.8)	(0.9)	(0.3)	(0.2)	(100.0)				
記載なし	427	488	1,926	6,289	1,804	20,657	25,000	16,529	2,049	5,444	2,712	965	370	276	84,936	9,710	94,646	9.20	0.77
(%)	(0.5)	(0.6)	(2.3)	(7.4)	(2.1)	(24.3)	(29.4)	(19.5)	(2.4)	(6.4)	(3.2)	(1.1)	(0.4)	(0.3)	(100.0)				
総計	874	1,266	4,971	16,290	4,676	53,902	65,236	42,404	5,227	13,277	6,453	2,165	769	539	218,049	10,922	228,971	9.18	0.74
(%)	(0.4)	(0.6)	(2.3)	(7.5)	(2.1)	(24.7)	(29.9)	(19.4)	(2.4)	(6.1)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
平均	110.26	111.52	107.39	105.26	104.82	105.31	106.58	107.54	107.74	106.85	107.65	105.74	105.11	105.53	106.43	106.67	106.43		
標準偏差	36.80	35.16	34.20	32.84	32.93	32.48	33.11	33.78	33.70	34.38	34.85	36.40	35.88	36.47	33.31	33.57	33.32		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査