

表 3411 透析前補正カルシウム濃度(mg/dL) PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満	2	1	8	24	6	41	61	50	4	20	15	4	3	2	241	8	249	9.29	0.91
(%)	(0.8)	(0.4)	(3.3)	(10.0)	(2.5)	(17.0)	(25.3)	(20.7)	(1.7)	(8.3)	(6.2)	(1.7)	(1.2)	(0.8)	(100.0)				
20～	1	2	4	21	3	56	100	60	11	23	10	2	2	2	297	4	301	9.29	0.74
(%)	(0.3)	(0.7)	(1.3)	(7.1)	(1.0)	(18.9)	(33.7)	(20.2)	(3.7)	(7.7)	(3.4)	(0.7)	(0.7)	(0.7)	(100.0)				
35～		1	2	16	10	59	93	60	7	14	5	1	1		269	5	274	9.21	0.61
(%)		(0.4)	(0.7)	(5.9)	(3.7)	(21.9)	(34.6)	(22.3)	(2.6)	(5.2)	(1.9)	(0.4)	(0.4)		(100.0)				
50～			7	26	8	102	139	102	11	21	17			1	437	7	444	9.25	0.64
(%)			(1.6)	(5.9)	(1.8)	(23.3)	(31.8)	(23.3)	(2.5)	(4.8)	(3.9)	(0.5)	(0.2)	(0.2)	(100.0)				
75～		1	4	21	11	83	147	74	2	30	5	3			381	10	391	9.21	0.61
(%)		(0.3)	(1.0)	(5.5)	(2.9)	(21.8)	(38.6)	(19.4)	(0.5)	(7.9)	(1.3)	(0.8)			(100.0)				
100～	1	2	5	26	7	85	91	65	3	31	7	2			325	6	331	9.17	0.67
(%)	(0.3)	(0.6)	(1.5)	(8.0)	(2.2)	(26.2)	(28.0)	(20.0)	(0.9)	(9.5)	(2.2)	(0.6)			(100.0)				
125～			3	23	6	64	67	45	2	11	5	1			227	7	234	9.12	0.63
(%)			(1.3)	(10.1)	(2.6)	(28.2)	(29.5)	(19.8)	(0.9)	(4.8)	(2.2)	(0.4)			(100.0)				
150～						1	2	1							4		4	9.25	0.31
(%)						(25.0)	(50.0)	(25.0)							(100.0)				
151～	1		1	18	4	48	75	36	5	12	6	2		2	210	5	215	9.22	0.72
(%)	(0.5)		(0.5)	(8.6)	(1.9)	(22.9)	(35.7)	(17.1)	(2.4)	(5.7)	(2.9)	(1.0)		(1.0)	(100.0)				
180～	1	1	5	11	2	41	33	29	1	7	2	1			134	3	137	9.07	0.74
(%)	(0.7)	(0.7)	(3.7)	(8.2)	(1.5)	(30.6)	(24.6)	(21.6)	(0.7)	(5.2)	(1.5)	(0.7)			(100.0)				
210～	1		3	13	2	18	32	18	1	3	1	2			94	4	98	9.08	0.73
(%)	(1.1)		(3.2)	(13.8)	(2.1)	(19.1)	(34.0)	(19.1)	(1.1)	(3.2)	(1.1)	(2.1)			(100.0)				
240～	1		1	3	1	20	31	17	2	9	1	1			87	1	88	9.23	0.70
(%)	(1.1)		(1.1)	(3.4)	(1.1)	(23.0)	(35.6)	(19.5)	(2.3)	(10.3)	(1.1)	(1.1)			(100.0)				
300～			1	7	3	13	7	6	3	2					42		42	8.95	0.67
(%)			(2.4)	(16.7)	(7.1)	(31.0)	(16.7)	(14.3)	(7.1)	(4.8)					(100.0)				
350～				2		9	6	8		4					29		29	9.25	0.59
(%)				(6.9)		(31.0)	(20.7)	(27.6)		(13.8)					(100.0)				
400～		1		5		2	7	6		2		1			24		24	9.15	0.84
(%)		(4.2)		(20.8)		(8.3)	(29.2)	(25.0)		(8.3)		(4.2)			(100.0)				
500～			2	2	1	12	11	12	1	3		1			45		45	9.18	0.69
(%)			(4.4)	(4.4)	(2.2)	(26.7)	(24.4)	(26.7)	(2.2)	(6.7)		(2.2)			(100.0)				
合計	8	9	46	218	64	654	902	589	53	192	74	23	7	7	2,846	60	2,906	9.21	0.69
(%)	(0.3)	(0.3)	(1.6)	(7.7)	(2.2)	(23.0)	(31.7)	(20.7)	(1.9)	(6.7)	(2.6)	(0.8)	(0.2)	(0.2)	(100.0)				
記載なし						2	2	4		1			2		11	20	31	9.81	1.03
(%)						(18.2)	(18.2)	(36.4)		(9.1)			(18.2)		(100.0)				
総計	8	9	46	218	64	656	904	593	53	193	74	23	9	7	2,857	80	2,937	9.21	0.70
(%)	(0.3)	(0.3)	(1.6)	(7.6)	(2.2)	(23.0)	(31.6)	(20.8)	(1.9)	(6.8)	(2.6)	(0.8)	(0.3)	(0.2)	(100.0)				
平均	126.38	116.44	120.13	120.81	108.22	117.83	108.49	112.45	99.89	116.78	73.07	140.09	24.57	68.29	112.09	97.78	111.79		
標準偏差	102.46	134.33	118.54	109.64	92.43	103.21	107.18	112.62	102.22	132.39	57.99	149.18	17.96	67.76	108.81	66.61	108.12		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査