

表 3412 透析前補正カルシウム濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）

透析前ヘモグロビン濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
5未満				2		8	4	10	1	1					26		26	9.22	0.56
(%)				(7.7)		(30.8)	(15.4)	(38.5)	(3.8)	(3.8)					(100.0)				
5～			1	2	1	4	13	3		1	3	1	1		30	1	31	9.34	0.93
(%)			(3.3)	(6.7)	(3.3)	(13.3)	(43.3)	(10.0)		(3.3)	(10.0)	(3.3)	(3.3)		(100.0)				
6～			4	9	4	12	34	23	5	8	5	2	2	2	110	4	114	9.38	0.88
(%)			(3.6)	(8.2)	(3.6)	(10.9)	(30.9)	(20.9)	(4.5)	(7.3)	(4.5)	(1.8)	(1.8)	(1.8)	(100.0)				
7～	4	5	7	27	5	85	131	95	13	30	23	10	4		439	6	445	9.31	0.85
(%)	(0.9)	(1.1)	(1.6)	(6.2)	(1.1)	(19.4)	(29.8)	(21.6)	(3.0)	(6.8)	(5.2)	(2.3)	(0.9)		(100.0)				
8～	8	8	37	109	33	380	447	295	32	135	76	26	13	8	1,607	30	1,637	9.28	0.83
(%)	(0.5)	(0.5)	(2.3)	(6.8)	(2.1)	(23.6)	(27.8)	(18.4)	(2.0)	(8.4)	(4.7)	(1.6)	(0.8)	(0.5)	(100.0)				
9～	13	49	157	461	127	1,341	1,692	1,088	132	378	217	80	22	20	5,777	73	5,850	9.20	0.78
(%)	(0.2)	(0.8)	(2.7)	(8.0)	(2.2)	(23.2)	(29.3)	(18.8)	(2.3)	(6.5)	(3.8)	(1.4)	(0.4)	(0.3)	(100.0)				
10～	41	72	318	971	287	3,175	3,825	2,550	290	821	340	96	38	18	12,842	105	12,947	9.17	0.72
(%)	(0.3)	(0.6)	(2.5)	(7.6)	(2.2)	(24.7)	(29.8)	(19.9)	(2.3)	(6.4)	(2.6)	(0.7)	(0.3)	(0.1)	(100.0)				
11～	54	73	258	838	237	2,848	3,523	2,269	277	666	304	98	30	15	11,490	87	11,577	9.17	0.73
(%)	(0.5)	(0.6)	(2.2)	(7.3)	(2.1)	(24.8)	(30.7)	(19.7)	(2.4)	(5.8)	(2.6)	(0.9)	(0.3)	(0.1)	(100.0)				
12～	23	35	109	315	83	1,033	1,280	876	101	286	125	45	15	5	4,331	51	4,382	9.19	0.75
(%)	(0.5)	(0.8)	(2.5)	(7.3)	(1.9)	(23.9)	(29.6)	(20.2)	(2.3)	(6.6)	(2.9)	(1.0)	(0.3)	(0.1)	(100.0)				
13～	10	13	31	94	25	240	350	244	28	74	50	14	4	1	1,178	7	1,185	9.21	0.78
(%)	(0.8)	(1.1)	(2.6)	(8.0)	(2.1)	(20.4)	(29.7)	(20.7)	(2.4)	(6.3)	(4.2)	(1.2)	(0.3)	(0.1)	(100.0)				
14～	7	3	21	28	10	122	146	93	16	38	20	9	2	2	517	6	523	9.21	0.86
(%)	(1.4)	(0.6)	(4.1)	(5.4)	(1.9)	(23.6)	(28.2)	(18.0)	(3.1)	(7.4)	(3.9)	(1.7)	(0.4)	(0.4)	(100.0)				
合計	160	258	943	2,856	812	9,248	11,445	7,546	895	2,438	1,163	381	131	71	38,347	370	38,717	9.19	0.75
(%)	(0.4)	(0.7)	(2.5)	(7.4)	(2.1)	(24.1)	(29.8)	(19.7)	(2.3)	(6.4)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
記載なし		1	8	12	6	53	70	66	8	14	7	2	1		248	981	1,229	9.24	0.66
(%)		(0.4)	(3.2)	(4.8)	(2.4)	(21.4)	(28.2)	(26.6)	(3.2)	(5.6)	(2.8)	(0.8)	(0.4)		(100.0)				
総計	160	259	951	2,868	818	9,301	11,515	7,612	903	2,452	1,170	383	132	71	38,595	1,351	39,946	9.19	0.75
(%)	(0.4)	(0.7)	(2.5)	(7.4)	(2.1)	(24.1)	(29.8)	(19.7)	(2.3)	(6.4)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
平均	11.23	10.92	10.87	10.83	10.81	10.85	10.85	10.85	10.86	10.80	10.73	10.69	10.50	10.35	10.84	10.62	10.84		
標準偏差	1.56	1.29	1.32	1.24	1.24	1.21	1.25	1.26	1.30	1.29	1.44	1.48	1.63	1.51	1.26	1.45	1.26		

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査