

表 3413 透析前補正カルシウム濃度(mg/dL) 総コレステロール濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

総コレステロール濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
100未満	1	6	24	79	17	220	275	198	28	74	32	15	8	5	982	10	992	9.26	0.78
(%)	(0.1)	(0.6)	(2.4)	(8.0)	(1.7)	(22.4)	(28.0)	(20.2)	(2.9)	(7.5)	(3.3)	(1.5)	(0.8)	(0.5)	(100.0)				
100～	14	20	56	235	82	718	871	575	59	186	95	40	12	5	2,968	20	2,988	9.19	0.76
(%)	(0.5)	(0.7)	(1.9)	(7.9)	(2.8)	(24.2)	(29.3)	(19.4)	(2.0)	(6.3)	(3.2)	(1.3)	(0.4)	(0.2)	(100.0)				
120～	22	34	132	432	117	1,397	1,735	1,042	135	338	192	49	20	8	5,653	41	5,694	9.18	0.73
(%)	(0.4)	(0.6)	(2.3)	(7.6)	(2.1)	(24.7)	(30.7)	(18.4)	(2.4)	(6.0)	(3.4)	(0.9)	(0.4)	(0.1)	(100.0)				
140～	26	39	157	495	145	1,691	2,053	1,400	147	418	211	55	25	6	6,868	37	6,905	9.18	0.73
(%)	(0.4)	(0.6)	(2.3)	(7.2)	(2.1)	(24.6)	(29.9)	(20.4)	(2.1)	(6.1)	(3.1)	(0.8)	(0.4)	(0.1)	(100.0)				
160～	28	51	169	437	122	1,401	1,739	1,223	162	410	156	55	16	10	5,979	32	6,011	9.18	0.76
(%)	(0.5)	(0.9)	(2.8)	(7.3)	(2.0)	(23.4)	(29.1)	(20.5)	(2.7)	(6.9)	(2.6)	(0.9)	(0.3)	(0.2)	(100.0)				
180～	21	24	114	271	68	894	1,174	786	93	227	114	40	16	4	3,846	27	3,873	9.19	0.75
(%)	(0.5)	(0.6)	(3.0)	(7.0)	(1.8)	(23.2)	(30.5)	(20.4)	(2.4)	(5.9)	(3.0)	(1.0)	(0.4)	(0.1)	(100.0)				
200～	8	13	42	169	29	493	710	454	44	176	67	19	6	4	2,234	19	2,253	9.23	0.71
(%)	(0.4)	(0.6)	(1.9)	(7.6)	(1.3)	(22.1)	(31.8)	(20.3)	(2.0)	(7.9)	(3.0)	(0.9)	(0.3)	(0.2)	(100.0)				
220～	3	7	29	86	27	195	272	210	22	60	25	14	2		952	4	956	9.18	0.74
(%)	(0.3)	(0.7)	(3.0)	(9.0)	(2.8)	(20.5)	(28.6)	(22.1)	(2.3)	(6.3)	(2.6)	(1.5)	(0.2)		(100.0)				
240～	3	3	10	23	4	70	111	92	12	37	12	4	1	1	383	1	384	9.28	0.83
(%)	(0.8)	(0.8)	(2.6)	(6.0)	(1.0)	(18.3)	(29.0)	(24.0)	(3.1)	(9.7)	(3.1)	(1.0)	(0.3)	(0.3)	(100.0)				
260～	2	2	9	7	3	42	70	52	6	8	8	1		3	213	3	216	9.25	0.87
(%)	(0.9)	(0.9)	(4.2)	(3.3)	(1.4)	(19.7)	(32.9)	(24.4)	(2.8)	(3.8)	(3.8)	(0.5)		(1.4)	(100.0)				
合計	128	199	742	2,234	614	7,121	9,010	6,032	708	1,934	912	292	106	46	30,078	194	30,272	9.19	0.74
(%)	(0.4)	(0.7)	(2.5)	(7.4)	(2.0)	(23.7)	(30.0)	(20.1)	(2.4)	(6.4)	(3.0)	(1.0)	(0.4)	(0.2)	(100.0)				
記載なし	32	60	209	634	204	2,180	2,505	1,580	195	518	258	91	26	25	8,517	1,157	9,674	9.17	0.75
(%)	(0.4)	(0.7)	(2.5)	(7.4)	(2.4)	(25.6)	(29.4)	(18.6)	(2.3)	(6.1)	(3.0)	(1.1)	(0.3)	(0.3)	(100.0)				
総計	160	259	951	2,868	818	9,301	11,515	7,612	903	2,452	1,170	383	132	71	38,595	1,351	39,946	9.19	0.75
(%)	(0.4)	(0.7)	(2.5)	(7.4)	(2.1)	(24.1)	(29.8)	(19.7)	(2.3)	(6.4)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
平均	162.22	160.04	161.15	157.14	154.96	156.67	158.26	159.46	158.69	159.01	156.10	156.06	149.60	159.65	158.01	156.63	158.00		
標準偏差	36.94	35.97	35.91	35.90	34.41	34.63	35.48	36.24	36.14	36.37	37.15	38.32	37.68	55.58	35.68	37.37	35.70		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査