

表 3415 透析前補正カルシウム濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
50未満	1	6	12	42	11	109	169	97	13	33	19	10	2	2	526	3	529	9.23	0.78
(%)	(0.2)	(1.1)	(2.3)	(8.0)	(2.1)	(20.7)	(32.1)	(18.4)	(2.5)	(6.3)	(3.6)	(1.9)	(0.4)	(0.4)	(100.0)				
50～	5	8	51	192	68	573	722	457	49	161	66	23	7		2,382	21	2,403	9.18	0.69
(%)	(0.2)	(0.3)	(2.1)	(8.1)	(2.9)	(24.1)	(30.3)	(19.2)	(2.1)	(6.8)	(2.8)	(1.0)	(0.3)		(100.0)				
70～	16	32	86	370	94	1,241	1,543	1,035	114	333	167	48	20	7	5,106	32	5,138	9.21	0.72
(%)	(0.3)	(0.6)	(1.7)	(7.2)	(1.8)	(24.3)	(30.2)	(20.3)	(2.2)	(6.5)	(3.3)	(0.9)	(0.4)	(0.1)	(100.0)				
90～	17	31	180	453	116	1,489	1,928	1,234	147	362	202	50	11	7	6,227	38	6,265	9.18	0.71
(%)	(0.3)	(0.5)	(2.9)	(7.3)	(1.9)	(23.9)	(31.0)	(19.8)	(2.4)	(5.8)	(3.2)	(0.8)	(0.2)	(0.1)	(100.0)				
110～	17	44	124	341	108	1,221	1,542	1,059	134	334	134	44	14	8	5,124	38	5,162	9.19	0.73
(%)	(0.3)	(0.9)	(2.4)	(6.7)	(2.1)	(23.8)	(30.1)	(20.7)	(2.6)	(6.5)	(2.6)	(0.9)	(0.3)	(0.2)	(100.0)				
130～	14	22	79	234	53	740	921	648	83	190	100	31	10	5	3,130	19	3,149	9.19	0.75
(%)	(0.4)	(0.7)	(2.5)	(7.5)	(1.7)	(23.6)	(29.4)	(20.7)	(2.7)	(6.1)	(3.2)	(1.0)	(0.3)	(0.2)	(100.0)				
150～	4	7	46	115	37	361	492	329	33	108	36	15	3	1	1,587	10	1,597	9.19	0.69
(%)	(0.3)	(0.4)	(2.9)	(7.2)	(2.3)	(22.7)	(31.0)	(20.7)	(2.1)	(6.8)	(2.3)	(0.9)	(0.2)	(0.1)	(100.0)				
170～	5	1	16	58	12	153	192	140	15	56	18	9	3		678	4	682	9.21	0.76
(%)	(0.7)	(0.1)	(2.4)	(8.6)	(1.8)	(22.6)	(28.3)	(20.6)	(2.2)	(8.3)	(2.7)	(1.3)	(0.4)		(100.0)				
190～	2	4	8	23	2	67	79	46	4	16	5	3	2		261	1	262	9.11	0.76
(%)	(0.8)	(1.5)	(3.1)	(8.8)	(0.8)	(25.7)	(30.3)	(17.6)	(1.5)	(6.1)	(1.9)	(1.1)	(0.8)		(100.0)				
210～	3	3	8	6	1	25	36	31	4	12	7			3	139	2	141	9.23	1.13
(%)	(2.2)	(2.2)	(5.8)	(4.3)	(0.7)	(18.0)	(25.9)	(22.3)	(2.9)	(8.6)	(5.0)			(2.2)	(100.0)				
合計	84	158	610	1,834	502	5,979	7,624	5,076	596	1,605	754	233	72	33	25,160	168	25,328	9.19	0.72
(%)	(0.3)	(0.6)	(2.4)	(7.3)	(2.0)	(23.8)	(30.3)	(20.2)	(2.4)	(6.4)	(3.0)	(0.9)	(0.3)	(0.1)	(100.0)				
記載なし	76	101	341	1,034	316	3,322	3,891	2,536	307	847	416	150	60	38	13,435	1,183	14,618	9.17	0.78
(%)	(0.6)	(0.8)	(2.5)	(7.7)	(2.4)	(24.7)	(29.0)	(18.9)	(2.3)	(6.3)	(3.1)	(1.1)	(0.4)	(0.3)	(100.0)				
総計	160	259	951	2,868	818	9,301	11,515	7,612	903	2,452	1,170	383	132	71	38,595	1,351	39,946	9.19	0.75
(%)	(0.4)	(0.7)	(2.5)	(7.4)	(2.1)	(24.1)	(29.8)	(19.7)	(2.3)	(6.4)	(3.0)	(1.0)	(0.3)	(0.2)	(100.0)				
平均	116.63	110.49	111.36	106.80	104.10	106.76	106.66	107.82	107.97	107.62	105.64	106.71	107.17	118.48	107.13	106.17	107.12		
標準偏差	39.64	33.97	35.07	34.35	32.78	32.61	33.11	33.42	32.57	34.05	34.37	35.20	34.64	49.87	33.37	33.64	33.37		

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査