

表 3424 補正カルシウム濃度(mg/dL) 体重(kg)別（腹膜透析患者）

体重	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
20未満						1	4	5	3	3	11	3	1	5	36	2	38	10.50	0.95
(%)						(2.8)	(11.1)	(13.9)	(8.3)	(8.3)	(30.6)	(8.3)	(2.8)	(13.9)	(100.0)				
20～			1	1			3	8	2	2	1	3			21	2	23	9.78	0.88
(%)			(4.8)	(4.8)			(14.3)	(38.1)	(9.5)	(9.5)	(4.8)	(14.3)			(100.0)				
30～		1	4	1		13	44	46	8	21	13	10	2	4	167	5	172	9.76	0.87
(%)		(0.6)	(2.4)	(0.6)		(7.8)	(26.3)	(27.5)	(4.8)	(12.6)	(7.8)	(6.0)	(1.2)	(2.4)	(100.0)				
40～	2	3	18	31	11	117	223	224	32	119	61	18	8		867	37	904	9.52	0.77
(%)	(0.2)	(0.3)	(2.1)	(3.6)	(1.3)	(13.5)	(25.7)	(25.8)	(3.7)	(13.7)	(7.0)	(2.1)	(0.9)		(100.0)				
50～	7	8	19	52	13	217	416	444	56	145	89	38	8	12	1,524	67	1,591	9.50	0.79
(%)	(0.5)	(0.5)	(1.2)	(3.4)	(0.9)	(14.2)	(27.3)	(29.1)	(3.7)	(9.5)	(5.8)	(2.5)	(0.5)	(0.8)	(100.0)				
60～	10	7	29	71	23	260	445	433	45	134	67	16	6	1	1,547	69	1,616	9.35	0.75
(%)	(0.6)	(0.5)	(1.9)	(4.6)	(1.5)	(16.8)	(28.8)	(28.0)	(2.9)	(8.7)	(4.3)	(1.0)	(0.4)	(0.1)	(100.0)				
70～	3	3	23	52	20	129	209	228	18	55	24	11	1	2	778	33	811	9.29	0.75
(%)	(0.4)	(0.4)	(3.0)	(6.7)	(2.6)	(16.6)	(26.9)	(29.3)	(2.3)	(7.1)	(3.1)	(1.4)	(0.1)	(0.3)	(100.0)				
80～	3	4	9	20	8	69	69	71	3	14	8	4		1	283	15	298	9.15	0.82
(%)	(1.1)	(1.4)	(3.2)	(7.1)	(2.8)	(24.4)	(24.4)	(25.1)	(1.1)	(4.9)	(2.8)	(1.4)		(0.4)	(100.0)				
90～		3	5	7	2	15	25	26	4	7	4	1	1		100	4	104	9.24	0.84
(%)		(3.0)	(5.0)	(7.0)	(2.0)	(15.0)	(25.0)	(26.0)	(4.0)	(7.0)	(4.0)	(1.0)	(1.0)		(100.0)				
100～				3		4	2	6		2	1				18	2	20	9.29	0.77
(%)				(16.7)		(22.2)	(11.1)	(33.3)		(11.1)	(5.6)				(100.0)				
110～			1	1	1	1	4	2							9	1	10	9.12	0.53
(%)			(11.1)	(11.1)	(11.1)	(11.1)	(44.4)	(22.2)							(100.0)				
120～			1			2		4	1	1					9		9	9.43	0.76
(%)			(11.1)			(22.2)		(44.4)	(11.1)	(11.1)					(100.0)				
合計	25	29	108	240	78	828	1,444	1,497	172	503	279	104	27	25	5,359	237	5,596	9.42	0.79
(%)	(0.5)	(0.5)	(2.0)	(4.5)	(1.5)	(15.5)	(26.9)	(27.9)	(3.2)	(9.4)	(5.2)	(1.9)	(0.5)	(0.5)	(100.0)				
記載なし	4	7	19	46	8	78	158	132	19	54	17	7	2	2	553	2,792	3,345	9.29	0.84
(%)	(0.7)	(1.3)	(3.4)	(8.3)	(1.4)	(14.1)	(28.6)	(23.9)	(3.4)	(9.8)	(3.1)	(1.3)	(0.4)	(0.4)	(100.0)				
総計	29	36	127	286	86	906	1,602	1,629	191	557	296	111	29	27	5,912	3,029	8,941	9.41	0.80
(%)	(0.5)	(0.6)	(2.1)	(4.8)	(1.5)	(15.3)	(27.1)	(27.6)	(3.2)	(9.4)	(5.0)	(1.9)	(0.5)	(0.5)	(100.0)				
平均	64.27	65.53	63.16	65.48	66.63	63.23	60.82	60.95	57.36	57.96	55.72	53.68	53.60	46.53	60.72	61.21	60.74		
標準偏差	11.39	16.16	14.71	14.24	13.09	13.39	12.87	13.61	14.54	13.83	14.94	15.38	14.07	20.04	13.88	14.27	13.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査