

表 3429 補正カルシウム濃度(mg/dL) 総コレステロール濃度(mg/dL)別（腹膜透析患者）

総コレステロール濃度	7.0未満	7.0～	7.5～	8.0～	8.4～	8.5～	9.0～	9.5～	10.0～	10.1～	10.5～	11.0～	11.5～	12.0～	合計	記載なし	総計	平均	標準偏差
100未満	1	1	1	4	1	2	12	11	2	5	7	2		1	50		50	9.56	1.03
(%)	(2.0)	(2.0)	(2.0)	(8.0)	(2.0)	(4.0)	(24.0)	(22.0)	(4.0)	(10.0)	(14.0)	(4.0)		(2.0)	(100.0)				
100～	2		2	11	4	26	45	38	6	14	5	3	1		157	3	160	9.31	0.83
(%)	(1.3)		(1.3)	(7.0)	(2.5)	(16.6)	(28.7)	(24.2)	(3.8)	(8.9)	(3.2)	(1.9)	(0.6)		(100.0)				
120～	4	2	11	22	10	71	114	117	15	44	21	6	2		439	9	448	9.36	0.80
(%)	(0.9)	(0.5)	(2.5)	(5.0)	(2.3)	(16.2)	(26.0)	(26.7)	(3.4)	(10.0)	(4.8)	(1.4)	(0.5)		(100.0)				
140～	3	8	8	43	7	135	220	215	19	71	43	11	1	3	787	19	806	9.38	0.78
(%)	(0.4)	(1.0)	(1.0)	(5.5)	(0.9)	(17.2)	(28.0)	(27.3)	(2.4)	(9.0)	(5.5)	(1.4)	(0.1)	(0.4)	(100.0)				
160～	3	3	19	31	12	130	233	239	28	87	38	24	4		851	14	865	9.43	0.76
(%)	(0.4)	(0.4)	(2.2)	(3.6)	(1.4)	(15.3)	(27.4)	(28.1)	(3.3)	(10.2)	(4.5)	(2.8)	(0.5)		(100.0)				
180～	2	4	16	40	13	111	227	236	24	84	44	10	2	5	818	15	833	9.42	0.77
(%)	(0.2)	(0.5)	(2.0)	(4.9)	(1.6)	(13.6)	(27.8)	(28.9)	(2.9)	(10.3)	(5.4)	(1.2)	(0.2)	(0.6)	(100.0)				
200～	3	3	9	31	5	79	147	162	17	58	27	7	5	1	554	10	564	9.42	0.77
(%)	(0.5)	(0.5)	(1.6)	(5.6)	(0.9)	(14.3)	(26.5)	(29.2)	(3.1)	(10.5)	(4.9)	(1.3)	(0.9)	(0.2)	(100.0)				
220～			7	23	7	36	86	90	11	30	18	5	1	1	315	3	318	9.43	0.75
(%)			(2.2)	(7.3)	(2.2)	(11.4)	(27.3)	(28.6)	(3.5)	(9.5)	(5.7)	(1.6)	(0.3)	(0.3)	(100.0)				
240～	1		1	2	2	21	54	40	8	12	9	4		3	157	2	159	9.54	0.80
(%)	(0.6)		(0.6)	(1.3)	(1.3)	(13.4)	(34.4)	(25.5)	(5.1)	(7.6)	(5.7)	(2.5)		(1.9)	(100.0)				
260～	1		4	9		19	32	41	4	15	12	3	2	4	146	3	149	9.57	0.94
(%)	(0.7)		(2.7)	(6.2)		(13.0)	(21.9)	(28.1)	(2.7)	(10.3)	(8.2)	(2.1)	(1.4)	(2.7)	(100.0)				
合計	20	21	78	216	61	630	1,170	1,189	134	420	224	75	18	18	4,274	78	4,352	9.42	0.78
(%)	(0.5)	(0.5)	(1.8)	(5.1)	(1.4)	(14.7)	(27.4)	(27.8)	(3.1)	(9.8)	(5.2)	(1.8)	(0.4)	(0.4)	(100.0)				
記載なし	9	15	49	70	25	276	432	440	57	137	72	36	11	9	1,638	2,951	4,589	9.37	0.82
(%)	(0.5)	(0.9)	(3.0)	(4.3)	(1.5)	(16.8)	(26.4)	(26.9)	(3.5)	(8.4)	(4.4)	(2.2)	(0.7)	(0.5)	(100.0)				
総計	29	36	127	286	86	906	1,602	1,629	191	557	296	111	29	27	5,912	3,029	8,941	9.41	0.80
(%)	(0.5)	(0.6)	(2.1)	(4.8)	(1.5)	(15.3)	(27.1)	(27.6)	(3.2)	(9.4)	(5.0)	(1.9)	(0.5)	(0.5)	(100.0)				
平均	165.05	164.00	181.27	177.87	172.80	175.49	179.53	180.29	179.49	179.46	181.25	178.67	186.83	213.94	179.10	174.33	179.01		
標準偏差	46.89	29.12	42.92	42.20	40.05	37.82	40.56	40.02	40.48	40.07	44.23	45.54	43.05	59.37	40.54	40.37	40.54		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査