

表 3438 透析前リン濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
施設血液透析	9,509	11,580	20,015	29,543	34,905	35,151	30,141	5,109	17,156	14,905	9,199	5,539	3,298	2,084	3,276	231,410	24,231	255,641	5.19	1.43
(%)	(4.1)	(5.0)	(8.6)	(12.8)	(15.1)	(15.2)	(13.0)	(2.2)	(7.4)	(6.4)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
Off-line HDF	248	268	461	704	840	945	762	114	437	382	250	168	93	67	111	5,850	465	6,315	5.27	1.50
(%)	(4.2)	(4.6)	(7.9)	(12.0)	(14.4)	(16.2)	(13.0)	(1.9)	(7.5)	(6.5)	(4.3)	(2.9)	(1.6)	(1.1)	(1.9)	(100.0)				
On-line HDF	886	1,304	2,406	3,897	4,907	5,207	4,575	838	2,811	2,452	1,530	949	571	362	578	33,273	2,817	36,090	5.37	1.41
(%)	(2.7)	(3.9)	(7.2)	(11.7)	(14.7)	(15.6)	(13.7)	(2.5)	(8.4)	(7.4)	(4.6)	(2.9)	(1.7)	(1.1)	(1.7)	(100.0)				
Push/Pull HDF	18	33	52	64	84	90	64	12	40	21	17	9	3	6	5	518	19	537	5.06	1.33
(%)	(3.5)	(6.4)	(10.0)	(12.4)	(16.2)	(17.4)	(12.4)	(2.3)	(7.7)	(4.1)	(3.3)	(1.7)	(0.6)	(1.2)	(1.0)	(100.0)				
バイオフィльтраーション	15	18	23	35	48	51	38	6	32	22	17	9	3		7	324	17	341	5.25	1.42
(%)	(4.6)	(5.6)	(7.1)	(10.8)	(14.8)	(15.7)	(11.7)	(1.9)	(9.9)	(6.8)	(5.2)	(2.8)	(0.9)		(2.2)	(100.0)				
血液濾過	6	3	7	11	10	16	14	1	8	5	3	3	1	1	2	91	1	92	5.19	1.59
(%)	(6.6)	(3.3)	(7.7)	(12.1)	(11.0)	(17.6)	(15.4)	(1.1)	(8.8)	(5.5)	(3.3)	(3.3)	(1.1)	(1.1)	(2.2)	(100.0)				
血液吸着透析 (リクセル)	42	76	134	205	283	252	238	38	126	77	39	26	7	11	7	1,561	62	1,623	5.09	1.18
(%)	(2.7)	(4.9)	(8.6)	(13.1)	(18.1)	(16.1)	(15.2)	(2.4)	(8.1)	(4.9)	(2.5)	(1.7)	(0.4)	(0.7)	(0.4)	(100.0)				
血液吸着透析 (その他)									1			2				3	4	7	7.07	0.84
(%)									(33.3)			(66.7)				(100.0)				
在宅血液透析	36	30	59	53	61	58	34	5	21	15	10	4	6	3	2	397	124	521	4.70	1.38
(%)	(9.1)	(7.6)	(14.9)	(13.4)	(15.4)	(14.6)	(8.6)	(1.3)	(5.3)	(3.8)	(2.5)	(1.0)	(1.5)	(0.8)	(0.5)	(100.0)				
腹膜透析 (CAPD)	80	153	312	445	511	523	432	82	243	196	94	61	30	16	46	3,224	1,797	5,021	5.15	1.32
(%)	(2.5)	(4.7)	(9.7)	(13.8)	(15.8)	(16.2)	(13.4)	(2.5)	(7.5)	(6.1)	(2.9)	(1.9)	(0.9)	(0.5)	(1.4)	(100.0)				
腹膜透析 (APD)	35	92	176	216	295	297	276	43	174	142	86	59	29	18	30	1,968	871	2,839	5.33	1.39
(%)	(1.8)	(4.7)	(8.9)	(11.0)	(15.0)	(15.1)	(14.0)	(2.2)	(8.8)	(7.2)	(4.4)	(3.0)	(1.5)	(0.9)	(1.5)	(100.0)				
腹膜透析 (CCPD)	11	25	37	109	115	128	131	13	60	61	32	29	12	4	13	780	301	1,081	5.43	1.33
(%)	(1.4)	(3.2)	(4.7)	(14.0)	(14.7)	(16.4)	(16.8)	(1.7)	(7.7)	(7.8)	(4.1)	(3.7)	(1.5)	(0.5)	(1.7)	(100.0)				
合計	10,886	13,582	23,682	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
記載なし																				
総計	10,886	13,582	23,682	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2014年調査