

表 3443 透析前リン濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,915	3,763	6,863	10,720	13,320	14,062	12,061	2,064	6,828	5,920	3,628	2,168	1,248	782	1,207	87,549	9,421	96,970	5.26	1.39
(%)	(3.3)	(4.3)	(7.8)	(12.2)	(15.2)	(16.1)	(13.8)	(2.4)	(7.8)	(6.8)	(4.1)	(2.5)	(1.4)	(0.9)	(1.4)	(100.0)				
慢性腎盂腎炎	118	121	237	366	403	415	376	63	211	193	115	56	30	27	51	2,782	260	3,042	5.22	1.45
(%)	(4.2)	(4.3)	(8.5)	(13.2)	(14.5)	(14.9)	(13.5)	(2.3)	(7.6)	(6.9)	(4.1)	(2.0)	(1.1)	(1.0)	(1.8)	(100.0)				
急速進行性糸球体腎炎	126	164	218	293	347	311	277	42	142	104	75	53	21	18	29	2,220	274	2,494	5.00	1.45
(%)	(5.7)	(7.4)	(9.8)	(13.2)	(15.6)	(14.0)	(12.5)	(1.9)	(6.4)	(4.7)	(3.4)	(2.4)	(0.9)	(0.8)	(1.3)	(100.0)				
妊娠腎/妊娠中毒症	41	44	121	186	227	235	230	36	118	95	65	43	25	8	21	1,495	119	1,614	5.32	1.38
(%)	(2.7)	(2.9)	(8.1)	(12.4)	(15.2)	(15.7)	(15.4)	(2.4)	(7.9)	(6.4)	(4.3)	(2.9)	(1.7)	(0.5)	(1.4)	(100.0)				
その他の分類不能の腎炎	40	69	88	150	196	172	184	27	95	79	44	36	25	18	29	1,252	121	1,373	5.33	1.52
(%)	(3.2)	(5.5)	(7.0)	(12.0)	(15.7)	(13.7)	(14.7)	(2.2)	(7.6)	(6.3)	(3.5)	(2.9)	(2.0)	(1.4)	(2.3)	(100.0)				
多発性囊胞腎	258	362	683	1,226	1,449	1,676	1,476	250	887	744	406	268	134	77	127	10,023	983	11,006	5.33	1.33
(%)	(2.6)	(3.6)	(6.8)	(12.2)	(14.5)	(16.7)	(14.7)	(2.5)	(8.8)	(7.4)	(4.1)	(2.7)	(1.3)	(0.8)	(1.3)	(100.0)				
腎硬化症	1,116	1,519	2,523	3,385	3,995	3,824	3,221	560	1,747	1,463	907	542	309	207	327	25,645	2,653	28,298	5.09	1.41
(%)	(4.4)	(5.9)	(9.8)	(13.2)	(15.6)	(14.9)	(12.6)	(2.2)	(6.8)	(5.7)	(3.5)	(2.1)	(1.2)	(0.8)	(1.3)	(100.0)				
悪性高血圧	83	105	201	287	304	346	266	46	176	158	126	86	50	29	50	2,313	260	2,573	5.37	1.55
(%)	(3.6)	(4.5)	(8.7)	(12.4)	(13.1)	(15.0)	(11.5)	(2.0)	(7.6)	(6.8)	(5.4)	(3.7)	(2.2)	(1.3)	(2.2)	(100.0)				
糖尿病性腎症	4,379	5,300	9,258	13,609	15,951	15,821	13,617	2,279	8,002	6,979	4,398	2,629	1,621	988	1,577	106,408	11,673	118,081	5.20	1.44
(%)	(4.1)	(5.0)	(8.7)	(12.8)	(15.0)	(14.9)	(12.8)	(2.1)	(7.5)	(6.6)	(4.1)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
SLE腎炎	77	80	149	249	261	331	277	52	155	123	84	46	51	26	40	2,001	235	2,236	5.33	1.49
(%)	(3.8)	(4.0)	(7.4)	(12.4)	(13.0)	(16.5)	(13.8)	(2.6)	(7.7)	(6.1)	(4.2)	(2.3)	(2.5)	(1.3)	(2.0)	(100.0)				
アミロイド腎	22	23	38	43	56	49	59	13	26	29	15	8	7	1	3	392	59	451	5.10	1.43
(%)	(5.6)	(5.9)	(9.7)	(11.0)	(14.3)	(12.5)	(15.1)	(3.3)	(6.6)	(7.4)	(3.8)	(2.0)	(1.8)	(0.3)	(0.8)	(100.0)				
痛風腎	28	47	79	123	154	156	117	17	83	83	41	31	17	16	21	1,013	100	1,113	5.37	1.48
(%)	(2.8)	(4.6)	(7.8)	(12.1)	(15.2)	(15.4)	(11.5)	(1.7)	(8.2)	(8.2)	(4.0)	(3.1)	(1.7)	(1.6)	(2.1)	(100.0)				
先天性代謝異常等による腎不全	9	8	18	18	39	36	34	5	17	20	15	6	5	2	9	241	31	272	5.51	1.63
(%)	(3.7)	(3.3)	(7.5)	(7.5)	(16.2)	(14.9)	(14.1)	(2.1)	(7.1)	(8.3)	(6.2)	(2.5)	(2.1)	(0.8)	(3.7)	(100.0)				
腎・尿路結核	5	13	18	20	30	32	17	8	8	14	8	5	4	1	1	184	22	206	5.16	1.36
(%)	(2.7)	(7.1)	(9.8)	(10.9)	(16.3)	(17.4)	(9.2)	(4.3)	(4.3)	(7.6)	(4.3)	(2.7)	(2.2)	(0.5)	(0.5)	(100.0)				
腎・尿路結石	18	31	59	56	69	83	75	17	36	26	21	18	5	5	6	525	56	581	5.17	1.43
(%)	(3.4)	(5.9)	(11.2)	(10.7)	(13.1)	(15.8)	(14.3)	(3.2)	(6.9)	(5.0)	(4.0)	(3.4)	(1.0)	(1.0)	(1.1)	(100.0)				
腎・尿路腫瘍	33	37	73	100	119	129	90	19	48	57	36	17	15	6	9	788	98	886	5.18	1.42
(%)	(4.2)	(4.7)	(9.3)	(12.7)	(15.1)	(16.4)	(11.4)	(2.4)	(6.1)	(7.2)	(4.6)	(2.2)	(1.9)	(0.8)	(1.1)	(100.0)				
閉塞性尿路障害	19	37	72	82	90	80	84	12	41	56	30	11	9	9	12	644	79	723	5.23	1.47
(%)	(3.0)	(5.7)	(11.2)	(12.7)	(14.0)	(12.4)	(13.0)	(1.9)	(6.4)	(8.7)	(4.7)	(1.7)	(1.4)	(1.4)	(1.9)	(100.0)				
骨髄腫	15	20	27	25	42	28	27	5	13	13	7	6	3		2	233	43	276	4.90	1.46
(%)	(6.4)	(8.6)	(11.6)	(10.7)	(18.0)	(12.0)	(11.6)	(2.1)	(5.6)	(5.6)	(3.0)	(2.6)	(1.3)		(0.9)	(100.0)				
腎形成不全	10	20	30	69	68	90	90	14	53	53	32	21	15	12	12	589	60	649	5.60	1.48
(%)	(1.7)	(3.4)	(5.1)	(11.7)	(11.5)	(15.3)	(15.3)	(2.4)	(9.0)	(9.0)	(5.4)	(3.6)	(2.5)	(2.0)	(2.0)	(100.0)				
不明	1,153	1,405	2,193	3,173	3,693	3,542	3,031	538	1,760	1,505	886	563	329	248	377	24,396	3,148	27,544	5.14	1.47
(%)	(4.7)	(5.8)	(9.0)	(13.0)	(15.1)	(14.5)	(12.4)	(2.2)	(7.2)	(6.2)	(3.6)	(2.3)	(1.3)	(1.0)	(1.5)	(100.0)				
再導入	65	63	127	209	269	302	278	51	171	150	94	62	39	36	43	1,959	204	2,163	5.48	1.46
(%)	(3.3)	(3.2)	(6.5)	(10.7)	(13.7)	(15.4)	(14.2)	(2.6)	(8.7)	(7.7)	(4.8)	(3.2)	(2.0)	(1.8)	(2.2)	(100.0)				
その他	356	351	607	893	977	998	818	143	492	414	244	183	91	56	124	6,747	810	7,557	5.15	1.49
(%)	(5.3)	(5.2)	(9.0)	(13.2)	(14.5)	(14.8)	(12.1)	(2.1)	(7.3)	(6.1)	(3.8)	(2.7)	(1.3)	(0.8)	(1.8)	(100.0)				
合計	10,886	13,582	23,882	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
記載なし																				
総計	10,886	13,582	23,882	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査