

表 3447 透析前リン濃度(mg/dL) PTH値(pg/mL)別 (透析患者全体) PTH測定法:whole-PTH

PTH値	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	136	144	213	245	299	283	204	38	106	92	54	35	17	5	19	1,890	4	1,894	4.84	1.41
(%)	(7.2)	(7.6)	(11.3)	(13.0)	(15.8)	(15.0)	(10.8)	(2.0)	(5.6)	(4.9)	(2.9)	(1.9)	(0.9)	(0.3)	(1.0)	(100.0)				
20～	128	180	273	317	393	380	244	45	126	114	54	33	20	6	15	2,328	3	2,331	4.84	1.31
(%)	(5.5)	(7.7)	(11.7)	(13.6)	(16.9)	(16.3)	(10.5)	(1.9)	(5.4)	(4.9)	(2.3)	(1.4)	(0.9)	(0.3)	(0.6)	(100.0)				
35～	95	131	232	305	334	325	232	27	136	91	50	36	16	14	10	2,034	9	2,043	4.91	1.29
(%)	(4.7)	(6.4)	(11.4)	(15.0)	(16.4)	(16.0)	(11.4)	(1.3)	(6.7)	(4.5)	(2.5)	(1.8)	(0.8)	(0.7)	(0.5)	(100.0)				
50～	138	174	342	436	528	496	415	60	182	199	80	41	21	11	19	3,142	11	3,153	4.97	1.27
(%)	(4.4)	(5.5)	(10.9)	(13.9)	(16.8)	(15.8)	(13.2)	(1.9)	(5.8)	(6.3)	(2.5)	(1.3)	(0.7)	(0.4)	(0.6)	(100.0)				
75～	67	138	256	391	467	436	355	65	194	151	91	47	30	21	29	2,738	10	2,748	5.12	1.29
(%)	(2.4)	(5.0)	(9.3)	(14.3)	(17.1)	(15.9)	(13.0)	(2.4)	(7.1)	(5.5)	(3.3)	(1.7)	(1.1)	(0.8)	(1.1)	(100.0)				
100～	68	106	198	275	377	358	292	70	173	159	83	57	31	15	20	2,282	16	2,298	5.21	1.33
(%)	(3.0)	(4.6)	(8.7)	(12.1)	(16.5)	(15.7)	(12.8)	(3.1)	(7.6)	(7.0)	(3.6)	(2.5)	(1.4)	(0.7)	(0.9)	(100.0)				
125～	36	72	114	193	220	269	218	31	134	95	66	36	18	8	17	1,527	12	1,539	5.26	1.30
(%)	(2.4)	(4.7)	(7.5)	(12.6)	(14.4)	(17.6)	(14.3)	(2.0)	(8.8)	(6.2)	(4.3)	(2.4)	(1.2)	(0.5)	(1.1)	(100.0)				
150～	1	2	4	3	5	10	12	1	5	2	1	1	1			48		48	5.26	1.15
(%)	(2.1)	(4.2)	(8.3)	(6.3)	(10.4)	(20.8)	(25.0)	(2.1)	(10.4)	(4.2)	(2.1)	(2.1)	(2.1)			(100.0)				
151～	21	40	86	139	187	217	204	38	118	96	51	32	14	7	9	1,259	4	1,263	5.36	1.24
(%)	(1.7)	(3.2)	(6.8)	(11.0)	(14.9)	(17.2)	(16.2)	(3.0)	(9.4)	(7.6)	(4.1)	(2.5)	(1.1)	(0.6)	(0.7)	(100.0)				
180～	18	17	51	89	148	142	128	23	87	64	34	32	23	6	9	871	4	875	5.46	1.30
(%)	(2.1)	(2.0)	(5.9)	(10.2)	(17.0)	(16.3)	(14.7)	(2.6)	(10.0)	(7.3)	(3.9)	(3.7)	(2.6)	(0.7)	(1.0)	(100.0)				
210～	9	25	35	73	99	94	91	15	54	46	37	14	10	6	19	627	1	628	5.48	1.42
(%)	(1.4)	(4.0)	(5.6)	(11.6)	(15.8)	(15.0)	(14.5)	(2.4)	(8.6)	(7.3)	(5.9)	(2.2)	(1.6)	(1.0)	(3.0)	(100.0)				
240～	15	21	38	65	96	107	107	15	54	53	39	30	13	4	15	672	2	674	5.53	1.40
(%)	(2.2)	(3.1)	(5.7)	(9.7)	(14.3)	(15.9)	(15.9)	(2.2)	(8.0)	(7.9)	(5.8)	(4.5)	(1.9)	(0.6)	(2.2)	(100.0)				
300～	2	7	12	21	40	47	49	3	24	29	21	11	3	8	7	284		284	5.77	1.43
(%)	(0.7)	(2.5)	(4.2)	(7.4)	(14.1)	(16.5)	(17.3)	(1.1)	(8.5)	(10.2)	(7.4)	(3.9)	(1.1)	(2.8)	(2.5)	(100.0)				
350～	2	3	9	10	25	26	32	5	17	16	12	9	4	2	5	177	2	179	5.81	1.43
(%)	(1.1)	(1.7)	(5.1)	(5.6)	(14.1)	(14.7)	(18.1)	(2.8)	(9.6)	(9.0)	(6.8)	(5.1)	(2.3)	(1.1)	(2.8)	(100.0)				
400～	4	9	7	17	23	22	30	5	24	17	17	8	2	6	10	201	1	202	5.85	1.62
(%)	(2.0)	(4.5)	(3.5)	(8.5)	(11.4)	(10.9)	(14.9)	(2.5)	(11.9)	(8.5)	(8.5)	(4.0)	(1.0)	(3.0)	(5.0)	(100.0)				
500～	2	10	12	24	25	34	20	11	28	16	16	13	9	5	22	247		247	6.08	1.99
(%)	(0.8)	(4.0)	(4.9)	(9.7)	(10.1)	(13.8)	(8.1)	(4.5)	(11.3)	(6.5)	(6.5)	(5.3)	(3.6)	(2.0)	(8.9)	(100.0)				
合計	742	1,079	1,882	2,603	3,266	3,246	2,633	452	1,462	1,240	706	435	232	124	225	20,327	79	20,406	5.13	1.35
(%)	(3.7)	(5.3)	(9.3)	(12.8)	(16.1)	(16.0)	(13.0)	(2.2)	(7.2)	(6.1)	(3.5)	(2.1)	(1.1)	(0.6)	(1.1)	(100.0)				
記載なし	29	34	40	54	43	46	29	3	17	13	11	9	3	1	1	333	324	657	4.70	1.40
(%)	(8.7)	(10.2)	(12.0)	(16.2)	(12.9)	(13.8)	(8.7)	(0.9)	(5.1)	(3.9)	(3.3)	(2.7)	(0.9)	(0.3)	(0.3)	(100.0)				
総計	771	1,113	1,922	2,657	3,309	3,292	2,662	455	1,479	1,253	717	444	235	125	226	20,660	403	21,063	5.12	1.35
(%)	(3.7)	(5.4)	(9.3)	(12.9)	(16.0)	(15.9)	(12.9)	(2.2)	(7.2)	(6.1)	(3.5)	(2.1)	(1.1)	(0.6)	(1.1)	(100.0)				
平均	72.15	85.09	87.13	96.70	101.39	105.94	116.86	122.48	125.35	120.64	140.51	148.22	144.79	160.87	204.50	107.99	111.43	108.01		
標準偏差	78.34	92.04	109.95	100.56	94.48	99.72	127.50	133.06	122.68	107.00	138.57	173.12	150.28	152.73	222.31	114.13	79.04	114.01		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査