

表 3448 透析前リン濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別 (透析患者全体)

透析前ヘモグロビン濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
5未満	18	14	25	39	52	49	37	6	16	19	10	11	2	1	3	302	1	303	5.05	1.39
(%)	(6.0)	(4.6)	(8.3)	(12.9)	(17.2)	(16.2)	(12.3)	(2.0)	(5.3)	(6.3)	(3.3)	(3.6)	(0.7)	(0.3)	(1.0)	(100.0)				
5～	34	26	32	46	65	61	47	5	29	32	21	7	8	4	9	426	3	429	5.12	1.63
(%)	(8.0)	(6.1)	(7.5)	(10.8)	(15.3)	(14.3)	(11.0)	(1.2)	(6.8)	(7.5)	(4.9)	(1.6)	(1.9)	(0.9)	(2.1)	(100.0)				
6～	165	93	123	126	140	132	117	20	57	55	36	19	17	17	19	1,136	24	1,160	4.73	1.77
(%)	(14.5)	(8.2)	(10.8)	(11.1)	(12.3)	(11.6)	(10.3)	(1.8)	(5.0)	(4.8)	(3.2)	(1.7)	(1.5)	(1.5)	(1.7)	(100.0)				
7～	526	375	500	566	572	501	424	61	263	189	115	104	63	36	68	4,363	72	4,435	4.75	1.67
(%)	(12.1)	(8.6)	(11.5)	(13.0)	(13.1)	(11.5)	(9.7)	(1.4)	(6.0)	(4.3)	(2.6)	(2.4)	(1.4)	(0.8)	(1.6)	(100.0)				
8～	1,338	1,123	1,625	1,975	2,012	1,822	1,485	268	822	673	482	275	196	144	180	14,420	134	14,554	4.84	1.55
(%)	(9.3)	(7.8)	(11.3)	(13.7)	(14.0)	(12.6)	(10.3)	(1.9)	(5.7)	(4.7)	(3.3)	(1.9)	(1.4)	(1.0)	(1.2)	(100.0)				
9～	2,486	2,820	4,567	6,328	6,947	6,696	5,348	912	2,986	2,567	1,576	1,022	571	374	591	45,791	198	45,989	5.04	1.44
(%)	(5.4)	(6.2)	(10.0)	(13.8)	(15.2)	(14.6)	(11.7)	(2.0)	(6.5)	(5.6)	(3.4)	(2.2)	(1.2)	(0.8)	(1.3)	(100.0)				
10～	3,069	4,424	8,083	12,366	14,834	14,972	12,439	2,133	7,043	5,774	3,486	2,026	1,183	722	1,127	93,681	231	93,912	5.18	1.36
(%)	(3.3)	(4.7)	(8.6)	(13.2)	(15.8)	(16.0)	(13.3)	(2.3)	(7.5)	(6.2)	(3.7)	(2.2)	(1.3)	(0.8)	(1.2)	(100.0)				
11～	2,072	3,128	5,995	9,364	11,953	12,344	11,060	1,832	6,316	5,671	3,326	2,023	1,100	712	1,109	78,005	168	78,173	5.31	1.37
(%)	(2.7)	(4.0)	(7.7)	(12.0)	(15.3)	(15.8)	(14.2)	(2.3)	(8.1)	(7.3)	(4.3)	(2.6)	(1.4)	(0.9)	(1.4)	(100.0)				
12～	765	1,035	1,935	3,208	3,883	4,443	4,069	730	2,495	2,289	1,527	884	614	349	550	28,776	78	28,854	5.46	1.44
(%)	(2.7)	(3.6)	(6.7)	(11.1)	(13.5)	(15.4)	(14.1)	(2.5)	(8.7)	(8.0)	(5.3)	(3.1)	(2.1)	(1.2)	(1.9)	(100.0)				
13～	202	295	428	684	924	981	1,007	170	668	670	440	282	164	124	246	7,285	23	7,308	5.63	1.58
(%)	(2.8)	(4.0)	(5.9)	(9.4)	(12.7)	(13.5)	(13.8)	(2.3)	(9.2)	(9.2)	(6.0)	(3.9)	(2.3)	(1.7)	(3.4)	(100.0)				
14～	80	98	145	237	305	334	342	65	243	205	177	139	99	63	134	2,666	7	2,673	5.81	1.72
(%)	(3.0)	(3.7)	(5.4)	(8.9)	(11.4)	(12.5)	(12.8)	(2.4)	(9.1)	(7.7)	(6.6)	(5.2)	(3.7)	(2.4)	(5.0)	(100.0)				
合計	10,755	13,431	23,458	34,939	41,687	42,335	36,375	6,202	20,938	18,144	11,196	6,792	4,017	2,546	4,036	276,851	939	277,790	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.6)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
記載なし	131	151	224	343	372	383	330	59	171	134	81	66	36	26	41	2,548	29,770	32,318	5.12	1.49
(%)	(5.1)	(5.9)	(8.8)	(13.5)	(14.6)	(15.0)	(13.0)	(2.3)	(6.7)	(5.3)	(3.2)	(2.6)	(1.4)	(1.0)	(1.6)	(100.0)				
総計	10,886	13,582	23,682	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
平均	10.16	10.42	10.52	10.62	10.69	10.75	10.81	10.82	10.85	10.90	10.92	10.93	10.94	10.93	11.00	10.71	10.06	10.71		
標準偏差	1.48	1.34	1.27	1.24	1.22	1.22	1.23	1.24	1.25	1.27	1.33	1.38	1.43	1.45	1.51	1.28	1.63	1.28		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査