

表 3451 透析前リン濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（透析患者全体）

Non-HDL-C濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
50未満	247	226	347	417	492	495	414	61	197	189	122	76	44	36	41	3,404	6	3,410	5.00	1.50
(%)	(7.3)	(6.6)	(10.2)	(12.3)	(14.5)	(14.5)	(12.2)	(1.8)	(5.8)	(5.6)	(3.6)	(2.2)	(1.3)	(1.1)	(1.2)	(100.0)				
50～	851	826	1,531	2,144	2,494	2,396	2,023	328	1,111	932	577	331	224	118	196	16,082	19	16,101	5.08	1.42
(%)	(5.3)	(5.1)	(9.5)	(13.3)	(15.5)	(14.9)	(12.6)	(2.0)	(6.9)	(5.8)	(3.6)	(2.1)	(1.4)	(0.7)	(1.2)	(100.0)				
70～	1,503	1,872	3,168	4,757	5,540	5,467	4,628	727	2,508	2,131	1,351	756	465	243	404	35,520	49	35,569	5.12	1.38
(%)	(4.2)	(5.3)	(8.9)	(13.4)	(15.6)	(15.4)	(13.0)	(2.0)	(7.1)	(6.0)	(3.8)	(2.1)	(1.3)	(0.7)	(1.1)	(100.0)				
90～	1,483	2,093	3,661	5,565	6,497	6,569	5,577	975	3,235	2,612	1,655	977	530	373	536	42,338	40	42,378	5.19	1.39
(%)	(3.5)	(4.9)	(8.6)	(13.1)	(15.3)	(15.5)	(13.2)	(2.3)	(7.6)	(6.2)	(3.9)	(2.3)	(1.3)	(0.9)	(1.3)	(100.0)				
110～	1,029	1,529	2,852	4,219	5,261	5,339	4,655	815	2,707	2,264	1,349	810	504	321	512	34,166	37	34,203	5.26	1.39
(%)	(3.0)	(4.5)	(8.3)	(12.3)	(15.4)	(15.6)	(13.6)	(2.4)	(7.9)	(6.6)	(3.9)	(2.4)	(1.5)	(0.9)	(1.5)	(100.0)				
130～	593	863	1,604	2,502	3,137	3,367	2,830	483	1,690	1,515	893	581	345	212	317	20,932	31	20,963	5.32	1.40
(%)	(2.8)	(4.1)	(7.7)	(12.0)	(15.0)	(16.1)	(13.5)	(2.3)	(8.1)	(7.2)	(4.3)	(2.8)	(1.6)	(1.0)	(1.5)	(100.0)				
150～	282	402	793	1,244	1,524	1,634	1,447	253	857	797	525	312	170	124	215	10,579	11	10,590	5.40	1.45
(%)	(2.7)	(3.8)	(7.5)	(11.8)	(14.4)	(15.4)	(13.7)	(2.4)	(8.1)	(7.5)	(5.0)	(2.9)	(1.6)	(1.2)	(2.0)	(100.0)				
170～	142	169	366	527	662	682	662	126	335	384	263	135	100	61	100	4,714	8	4,722	5.43	1.50
(%)	(3.0)	(3.6)	(7.8)	(11.2)	(14.0)	(14.5)	(14.0)	(2.7)	(7.1)	(8.1)	(5.6)	(2.9)	(2.1)	(1.3)	(2.1)	(100.0)				
190～	50	68	144	192	251	255	250	44	148	144	92	61	41	18	49	1,807	2	1,809	5.46	1.52
(%)	(2.8)	(3.8)	(8.0)	(10.6)	(13.9)	(14.1)	(13.8)	(2.4)	(8.2)	(8.0)	(5.1)	(3.4)	(2.3)	(1.0)	(2.7)	(100.0)				
210～	35	54	86	95	146	164	150	16	92	87	61	47	24	20	42	1,119	5	1,124	5.57	1.68
(%)	(3.1)	(4.8)	(7.7)	(8.5)	(13.0)	(14.7)	(13.4)	(1.4)	(8.2)	(7.8)	(5.5)	(4.2)	(2.1)	(1.8)	(3.8)	(100.0)				
合計	6,215	8,102	14,552	21,662	26,004	26,368	22,636	3,828	12,880	11,055	6,888	4,086	2,447	1,526	2,412	170,661	208	170,869	5.22	1.41
(%)	(3.6)	(4.7)	(8.5)	(12.7)	(15.2)	(15.5)	(13.3)	(2.2)	(7.5)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	4,671	5,480	9,130	13,620	16,055	16,350	14,069	2,433	8,229	7,223	4,389	2,772	1,606	1,046	1,665	108,738	30,501	139,239	5.21	1.45
(%)	(4.3)	(5.0)	(8.4)	(12.5)	(14.8)	(15.0)	(12.9)	(2.2)	(7.6)	(6.6)	(4.0)	(2.5)	(1.5)	(1.0)	(1.5)	(100.0)				
総計	10,886	13,582	23,682	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
平均	100.03	103.33	104.87	105.29	106.31	107.27	107.99	108.98	108.77	110.19	110.44	111.37	110.99	112.71	114.13	107.08	108.97	107.08		
標準偏差	33.97	32.80	33.46	32.45	32.92	33.12	33.46	33.03	33.42	34.52	35.22	35.75	36.10	35.70	38.08	33.60	39.46	33.61		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査