

表 3452 透析前リン濃度(mg/dL) 透析前収縮期血圧(mmHg)別 (透析患者全体)

透析前収縮期血圧	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
60未満	4	4	5	4	2	4	3			3	3	3		2	2	39	1	40	5.25	2.15
(%)	(10.3)	(10.3)	(12.8)	(10.3)	(5.1)	(10.3)	(7.7)			(7.7)	(7.7)	(7.7)		(5.1)	(5.1)	(100.0)				
60～	30	26	31	51	60	57	44	12	27	28	18	13	9	3	14	423	10	433	5.20	1.71
(%)	(7.1)	(6.1)	(7.3)	(12.1)	(14.2)	(13.5)	(10.4)	(2.8)	(6.4)	(6.6)	(4.3)	(3.1)	(2.1)	(0.7)	(3.3)	(100.0)				
80～	247	211	304	428	445	471	379	71	201	184	120	72	50	33	53	3,269	69	3,338	5.02	1.56
(%)	(7.6)	(6.5)	(9.3)	(13.1)	(13.6)	(14.4)	(11.6)	(2.2)	(6.1)	(5.6)	(3.7)	(2.2)	(1.5)	(1.0)	(1.6)	(100.0)				
100～	959	1,000	1,611	2,278	2,599	2,526	2,134	384	1,270	1,025	643	366	219	125	217	17,356	251	17,607	5.08	1.45
(%)	(5.5)	(5.8)	(9.3)	(13.1)	(15.0)	(14.6)	(12.3)	(2.2)	(7.3)	(5.9)	(3.7)	(2.1)	(1.3)	(0.7)	(1.3)	(100.0)				
120～	2,141	2,773	4,690	6,832	8,033	8,211	6,923	1,212	3,845	3,235	1,918	1,198	699	404	626	52,740	531	53,271	5.14	1.39
(%)	(4.1)	(5.3)	(8.9)	(13.0)	(15.2)	(15.6)	(13.1)	(2.3)	(7.3)	(6.1)	(3.6)	(2.3)	(1.3)	(0.8)	(1.2)	(100.0)				
140～	2,902	3,848	6,805	10,221	12,319	12,552	10,796	1,764	6,081	5,137	3,122	1,846	1,061	681	1,034	80,169	653	80,822	5.20	1.38
(%)	(3.6)	(4.8)	(8.5)	(12.7)	(15.4)	(15.7)	(13.5)	(2.2)	(7.6)	(6.4)	(3.9)	(2.3)	(1.3)	(0.8)	(1.3)	(100.0)				
160～	1,942	2,507	4,766	7,197	8,682	8,884	7,690	1,317	4,465	4,036	2,510	1,485	904	529	877	57,791	404	58,195	5.27	1.42
(%)	(3.4)	(4.3)	(8.2)	(12.5)	(15.0)	(15.4)	(13.3)	(2.3)	(7.7)	(7.0)	(4.3)	(2.6)	(1.6)	(0.9)	(1.5)	(100.0)				
180～	737	1,006	1,678	2,634	3,149	3,259	2,894	503	1,769	1,651	1,019	645	389	279	467	22,079	145	22,224	5.36	1.50
(%)	(3.3)	(4.6)	(7.6)	(11.9)	(14.3)	(14.8)	(13.1)	(2.3)	(8.0)	(7.5)	(4.6)	(2.9)	(1.8)	(1.3)	(2.1)	(100.0)				
200～	162	228	395	676	741	817	751	128	425	384	303	164	117	73	142	5,506	40	5,546	5.44	1.53
(%)	(2.9)	(4.1)	(7.2)	(12.3)	(13.5)	(14.8)	(13.6)	(2.3)	(7.7)	(7.0)	(5.5)	(3.0)	(2.1)	(1.3)	(2.6)	(100.0)				
220～	40	35	89	114	159	164	148	18	86	92	63	39	34	23	44	1,148	12	1,160	5.56	1.65
(%)	(3.5)	(3.0)	(7.8)	(9.9)	(13.9)	(14.3)	(12.9)	(1.6)	(7.5)	(8.0)	(5.5)	(3.4)	(3.0)	(2.0)	(3.8)	(100.0)				
合計	9,164	11,638	20,374	30,435	36,189	36,945	31,762	5,409	18,169	15,775	9,719	5,831	3,482	2,152	3,476	240,520	2,116	242,636	5.21	1.42
(%)	(3.8)	(4.8)	(8.5)	(12.7)	(15.0)	(15.4)	(13.2)	(2.2)	(7.6)	(6.6)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	1,722	1,944	3,308	4,847	5,870	5,773	4,943	852	2,940	2,503	1,558	1,027	571	420	601	38,879	28,593	67,472	5.21	1.46
(%)	(4.4)	(5.0)	(8.5)	(12.5)	(15.1)	(14.8)	(12.7)	(2.2)	(7.6)	(6.4)	(4.0)	(2.6)	(1.5)	(1.1)	(1.5)	(100.0)				
総計	10,886	13,582	23,682	35,282	42,059	42,718	36,705	6,261	21,109	18,278	11,277	6,858	4,053	2,572	4,077	279,399	30,709	310,108	5.21	1.43
(%)	(3.9)	(4.9)	(8.5)	(12.6)	(15.1)	(15.3)	(13.1)	(2.2)	(7.6)	(6.5)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
平均	147.25	148.84	149.93	150.54	150.85	151.11	151.61	151.21	151.97	152.80	153.32	153.36	154.21	155.14	155.76	151.18	145.39	151.13		
標準偏差	25.50	24.41	24.05	23.94	23.70	23.77	23.81	24.15	24.08	24.35	24.93	24.89	25.86	25.88	27.34	24.20	26.19	24.23		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査