

表 3458 透析前リン濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	2,332	2,950	5,324	8,186	10,051	10,626	9,153	1,534	5,125	4,465	2,755	1,616	935	586	893	66,531	2,046	68,577	5.24	1.39
(%)	(3.5)	(4.4)	(8.0)	(12.3)	(15.1)	(16.0)	(13.8)	(2.3)	(7.7)	(6.7)	(4.1)	(2.4)	(1.4)	(0.9)	(1.3)	(100.0)				
慢性腎盂腎炎	91	96	181	269	319	314	291	47	151	141	93	47	23	17	35	2,115	50	2,165	5.21	1.45
(%)	(4.3)	(4.5)	(8.6)	(12.7)	(15.1)	(14.8)	(13.8)	(2.2)	(7.1)	(6.7)	(4.4)	(2.2)	(1.1)	(0.8)	(1.7)	(100.0)				
急速進行性糸球体腎炎	108	130	176	232	277	238	229	27	115	84	57	43	16	13	20	1,765	69	1,834	4.97	1.44
(%)	(6.1)	(7.4)	(10.0)	(13.1)	(15.7)	(13.5)	(13.0)	(1.5)	(6.5)	(4.8)	(3.2)	(2.4)	(0.9)	(0.7)	(1.1)	(100.0)				
妊娠腎/妊娠中毒症	31	24	86	127	147	162	157	23	85	66	46	37	22	4	16	1,033	31	1,064	5.36	1.41
(%)	(3.0)	(2.3)	(8.3)	(12.3)	(14.2)	(15.7)	(15.2)	(2.2)	(8.2)	(6.4)	(4.5)	(3.6)	(2.1)	(0.4)	(1.5)	(100.0)				
その他の分類不能の腎炎	31	52	62	113	147	119	145	20	60	62	30	30	16	13	18	918	31	949	5.31	1.51
(%)	(3.4)	(5.7)	(6.8)	(12.3)	(16.0)	(13.0)	(15.8)	(2.2)	(6.5)	(6.8)	(3.3)	(3.3)	(1.7)	(1.4)	(2.0)	(100.0)				
多発性囊胞腎	217	293	552	994	1,157	1,332	1,185	200	692	582	313	200	105	58	99	7,979	219	8,198	5.31	1.32
(%)	(2.7)	(3.7)	(8.9)	(12.5)	(14.5)	(16.7)	(14.9)	(2.5)	(8.7)	(7.3)	(3.9)	(2.5)	(1.3)	(0.7)	(1.2)	(100.0)				
腎硬化症	910	1,251	2,076	2,744	3,266	3,131	2,619	456	1,385	1,186	724	435	254	170	265	20,872	588	21,460	5.08	1.41
(%)	(4.4)	(6.0)	(9.9)	(13.1)	(15.6)	(15.0)	(12.5)	(2.2)	(6.6)	(5.7)	(3.5)	(2.1)	(1.2)	(0.8)	(1.3)	(100.0)				
悪性高血圧	72	86	161	223	238	269	207	36	145	123	99	65	38	20	37	1,819	62	1,881	5.34	1.55
(%)	(4.0)	(4.7)	(8.9)	(12.3)	(13.1)	(14.8)	(11.4)	(2.0)	(8.0)	(6.8)	(5.4)	(3.6)	(2.1)	(1.1)	(2.0)	(100.0)				
糖尿病性腎症	3,705	4,424	7,631	11,279	13,166	13,069	11,188	1,889	6,528	5,708	3,569	2,126	1,335	809	1,259	87,685	2,822	90,507	5.19	1.44
(%)	(4.2)	(5.0)	(8.7)	(12.9)	(15.0)	(14.9)	(12.8)	(2.2)	(7.4)	(6.5)	(4.1)	(2.4)	(1.5)	(0.9)	(1.4)	(100.0)				
SLE腎炎	59	64	114	198	211	231	194	34	127	86	65	36	34	25	25	1,503	47	1,550	5.29	1.47
(%)	(3.9)	(4.3)	(7.6)	(13.2)	(14.0)	(15.4)	(12.9)	(2.3)	(8.4)	(5.7)	(4.3)	(2.4)	(2.3)	(1.7)	(1.7)	(100.0)				
アミロイド腎	21	18	27	33	43	36	50	9	22	22	10	4	5	1	3	304	21	325	5.07	1.46
(%)	(6.9)	(5.9)	(8.9)	(10.9)	(14.1)	(11.8)	(16.4)	(3.0)	(7.2)	(7.2)	(3.3)	(1.3)	(1.6)	(0.3)	(1.0)	(100.0)				
痛風腎	22	38	59	98	127	131	89	12	61	69	33	22	15	12	16	804	24	828	5.36	1.47
(%)	(2.7)	(4.7)	(7.3)	(12.2)	(15.8)	(16.3)	(11.1)	(1.5)	(7.6)	(8.6)	(4.1)	(2.7)	(1.9)	(1.5)	(2.0)	(100.0)				
先天性代謝異常 による腎不全	7	5	12	10	33	25	24	3	14	13	10	5	4	2	6	173	4	177	5.51	1.59
(%)	(4.0)	(2.9)	(6.9)	(5.8)	(19.1)	(14.5)	(13.9)	(1.7)	(8.1)	(7.5)	(5.8)	(2.9)	(2.3)	(1.2)	(3.5)	(100.0)				
腎・尿路結核	4	10	15	14	25	26	15	7	8	14	7	3	3	1	1	153	5	158	5.22	1.36
(%)	(2.6)	(6.5)	(9.8)	(9.2)	(16.3)	(17.0)	(9.8)	(4.6)	(5.2)	(9.2)	(4.6)	(2.0)	(2.0)	(0.7)	(0.7)	(100.0)				
腎・尿路結石	15	26	44	47	55	64	58	14	30	18	16	16	4	5	4	416	14	430	5.16	1.43
(%)	(3.6)	(6.3)	(10.6)	(11.3)	(13.2)	(15.4)	(13.9)	(3.4)	(7.2)	(4.3)	(3.8)	(3.8)	(1.0)	(1.2)	(1.0)	(100.0)				
腎・尿路腫瘍	29	34	61	85	99	108	73	15	43	45	32	13	10	5	8	660	26	686	5.15	1.42
(%)	(4.4)	(5.2)	(9.2)	(12.9)	(15.0)	(16.4)	(11.1)	(2.3)	(6.5)	(6.8)	(4.8)	(2.0)	(1.5)	(0.8)	(1.2)	(100.0)				
閉塞性尿路障害	14	29	59	67	67	60	70	10	29	41	28	6	6	6	10	502	15	517	5.21	1.47
(%)	(2.8)	(5.8)	(11.8)	(13.3)	(13.3)	(12.0)	(13.9)	(2.0)	(5.8)	(8.2)	(5.6)	(1.2)	(1.2)	(1.2)	(2.0)	(100.0)				
骨髄腫	10	13	19	19	29	22	21	4	12	11	7	5	3		2	177	13	190	5.04	1.51
(%)	(5.6)	(7.3)	(10.7)	(10.7)	(16.4)	(12.4)	(11.9)	(2.3)	(6.8)	(6.2)	(4.0)	(2.8)	(1.7)		(1.1)	(100.0)				
腎形成不全	9	15	19	49	48	65	54	10	39	36	21	18	9	9	10	411	8	419	5.59	1.53
(%)	(2.2)	(3.6)	(4.6)	(11.9)	(11.7)	(15.8)	(13.1)	(2.4)	(9.5)	(8.8)	(5.1)	(4.4)	(2.2)	(2.2)	(2.4)	(100.0)				
不明	934	1,100	1,763	2,549	2,969	2,779	2,365	427	1,338	1,169	701	454	261	193	299	19,301	824	20,125	5.13	1.47
(%)	(4.8)	(5.7)	(9.1)	(13.2)	(15.4)	(14.4)	(12.3)	(2.2)	(6.9)	(6.1)	(3.6)	(2.4)	(1.4)	(1.0)	(1.5)	(100.0)				
再導入	51	50	96	146	190	218	183	40	115	101	66	43	28	26	33	1,386	31	1,417	5.46	1.50
(%)	(3.7)	(3.6)	(6.9)	(10.5)	(13.7)	(15.7)	(13.2)	(2.9)	(8.3)	(7.3)	(4.8)	(3.1)	(2.0)	(1.9)	(2.4)	(100.0)				
その他	289	272	481	710	804	784	642	111	381	335	178	135	66	48	97	5,333	181	5,514	5.13	1.49
(%)	(5.4)	(5.1)	(9.0)	(13.3)	(15.1)	(14.7)	(12.0)	(2.1)	(7.1)	(6.3)	(3.3)	(2.5)	(1.2)	(0.9)	(1.8)	(100.0)				
合計	8,961	10,980	19,018	28,192	33,468	33,808	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし																				
総計	8,961	10,980	19,018	28,192	33,468	33,808	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査