

表 3460 透析前リン濃度(mg/dL) 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	6	4	9	13	12	16	13		4	3	2			1	1	84		84	4.83	1.35
(%)	(7.1)	(4.8)	(10.7)	(15.5)	(14.3)	(19.0)	(15.5)		(4.8)	(3.6)	(2.4)			(1.2)	(1.2)	(100.0)				
20～	104	84	90	107	110	100	64	6	37	27	17	14	10	5	10	785	17	802	4.57	1.58
(%)	(13.2)	(10.7)	(11.5)	(13.6)	(14.0)	(12.7)	(8.2)	(0.8)	(4.7)	(3.4)	(2.2)	(1.8)	(1.3)	(0.6)	(1.3)	(100.0)				
30～	1,615	1,396	2,065	2,653	2,806	2,606	2,045	349	1,094	905	524	343	167	112	196	18,876	241	19,117	4.82	1.47
(%)	(8.6)	(7.4)	(10.9)	(14.1)	(14.9)	(13.8)	(10.8)	(1.8)	(5.8)	(4.8)	(2.8)	(1.8)	(0.9)	(0.6)	(1.0)	(100.0)				
40～	3,190	3,480	5,795	8,273	9,190	8,936	7,411	1,166	3,930	3,195	1,900	1,121	631	392	540	59,150	659	59,809	5.00	1.38
(%)	(5.4)	(5.9)	(9.8)	(14.0)	(15.5)	(15.1)	(12.5)	(2.0)	(6.6)	(5.4)	(3.2)	(1.9)	(1.1)	(0.7)	(0.9)	(100.0)				
50～	2,479	3,585	6,162	9,344	11,163	11,124	9,420	1,665	5,273	4,610	2,787	1,530	861	557	846	71,406	703	72,109	5.18	1.37
(%)	(3.5)	(5.0)	(8.6)	(13.1)	(15.6)	(15.6)	(13.2)	(2.3)	(7.4)	(6.5)	(3.9)	(2.1)	(1.2)	(0.8)	(1.2)	(100.0)				
60～	1,036	1,653	3,257	5,155	6,455	6,961	6,162	1,070	3,583	3,230	1,931	1,235	764	475	723	43,690	392	44,082	5.37	1.40
(%)	(2.4)	(3.8)	(7.5)	(11.8)	(14.8)	(15.9)	(14.1)	(2.4)	(8.2)	(7.4)	(4.4)	(2.8)	(1.7)	(1.1)	(1.7)	(100.0)				
70～	309	511	1,105	1,699	2,414	2,579	2,391	402	1,550	1,454	989	606	398	245	381	17,033	151	17,184	5.57	1.45
(%)	(1.8)	(3.0)	(6.5)	(10.0)	(14.2)	(15.1)	(14.0)	(2.4)	(9.1)	(8.5)	(5.8)	(3.6)	(2.3)	(1.4)	(2.2)	(100.0)				
80～	109	121	278	551	754	858	839	157	619	536	392	286	201	113	214	6,028	53	6,081	5.81	1.55
(%)	(1.8)	(2.0)	(4.6)	(9.1)	(12.5)	(14.2)	(13.9)	(2.6)	(10.3)	(8.9)	(6.5)	(4.7)	(3.3)	(1.9)	(3.6)	(100.0)				
90～	26	46	81	158	247	273	328	53	214	217	159	122	72	56	122	2,174	21	2,195	6.04	1.63
(%)	(1.2)	(2.1)	(3.7)	(7.3)	(11.4)	(12.6)	(15.1)	(2.4)	(9.8)	(10.0)	(7.3)	(5.6)	(3.3)	(2.6)	(5.6)	(100.0)				
100～	7	14	35	42	60	95	117	23	68	77	59	40	31	32	48	748	7	755	6.20	1.66
(%)	(0.9)	(1.9)	(4.7)	(5.6)	(8.0)	(12.7)	(15.6)	(3.1)	(9.1)	(10.3)	(7.9)	(5.3)	(4.1)	(4.3)	(6.4)	(100.0)				
110～	3	4	9	24	27	31	38	3	29	33	19	11	23	13	29	296	3	299	6.40	1.94
(%)	(1.0)	(1.4)	(3.0)	(8.1)	(9.1)	(10.5)	(12.8)	(1.0)	(9.8)	(11.1)	(6.4)	(3.7)	(7.8)	(4.4)	(9.8)	(100.0)				
120～	2	3	5	8	13	15	28	3	17	17	22	7	17	12	19	188	1	189	6.63	1.82
(%)	(1.1)	(1.6)	(2.7)	(4.3)	(6.9)	(8.0)	(14.9)	(1.6)	(9.0)	(9.0)	(11.7)	(3.7)	(9.0)	(6.4)	(10.1)	(100.0)				
合計	8,886	10,901	18,891	28,027	33,251	33,594	28,856	4,897	16,418	14,304	8,801	5,315	3,175	2,013	3,129	220,458	2,248	222,706	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	75	79	127	165	217	215	156	31	87	73	59	44	17	10	27	1,382	4,883	6,265	5.13	1.54
(%)	(5.4)	(5.7)	(9.2)	(11.9)	(15.7)	(15.6)	(11.3)	(2.2)	(6.3)	(5.3)	(4.3)	(3.2)	(1.2)	(0.7)	(2.0)	(100.0)				
総計	8,961	10,980	19,018	28,192	33,468	33,809	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
平均	49.75	52.00	53.15	53.95	54.91	55.53	56.33	56.63	57.25	57.87	58.65	59.18	60.79	61.04	61.88	55.46	53.97	55.45		
標準偏差	11.62	11.41	11.70	11.85	12.16	12.41	12.93	12.70	13.27	13.57	14.18	14.54	15.64	16.26	16.97	12.90	12.77	12.90		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査