

表 3462 透析前リン濃度(mg/dL) PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	114	114	164	199	235	232	160	27	82	71	39	24	14	4	17	1,496	3	1,499	4.82	1.42
(%)	(7.6)	(7.6)	(11.0)	(13.3)	(15.7)	(15.5)	(10.7)	(1.8)	(5.5)	(4.7)	(2.6)	(1.6)	(0.9)	(0.3)	(1.1)	(100.0)				
20～	115	144	219	259	304	289	195	36	101	85	37	26	16	5	14	1,845	3	1,848	4.81	1.33
(%)	(6.2)	(7.8)	(11.9)	(14.0)	(16.5)	(15.7)	(10.6)	(2.0)	(5.5)	(4.6)	(2.0)	(1.4)	(0.9)	(0.3)	(0.8)	(100.0)				
35～	79	112	185	240	256	258	175	23	104	76	38	27	14	8	10	1,605	8	1,613	4.89	1.31
(%)	(4.9)	(7.0)	(11.5)	(15.0)	(16.0)	(16.1)	(10.9)	(1.4)	(6.5)	(4.7)	(2.4)	(1.7)	(0.9)	(0.5)	(0.6)	(100.0)				
50～	116	135	271	325	394	384	330	46	147	163	70	32	16	8	14	2,451	8	2,459	4.98	1.28
(%)	(4.7)	(5.5)	(11.1)	(13.3)	(16.1)	(15.7)	(13.5)	(1.9)	(6.0)	(6.7)	(2.9)	(1.3)	(0.7)	(0.3)	(0.6)	(100.0)				
75～	57	111	198	297	360	356	275	45	158	114	72	37	23	19	19	2,141	7	2,148	5.11	1.28
(%)	(2.7)	(5.2)	(9.2)	(13.9)	(16.8)	(16.6)	(12.8)	(2.1)	(7.4)	(5.3)	(3.4)	(1.7)	(1.1)	(0.9)	(0.9)	(100.0)				
100～	52	87	165	207	288	291	228	52	127	125	58	42	24	11	16	1,773	14	1,787	5.19	1.33
(%)	(2.9)	(4.9)	(9.3)	(11.7)	(16.2)	(16.4)	(12.9)	(2.9)	(7.2)	(7.1)	(3.3)	(2.4)	(1.4)	(0.6)	(0.9)	(100.0)				
125～	28	53	95	142	174	202	172	26	102	72	48	27	14	8	13	1,176	11	1,187	5.25	1.29
(%)	(2.4)	(4.5)	(8.1)	(12.1)	(14.8)	(17.2)	(14.6)	(2.2)	(8.7)	(6.1)	(4.1)	(2.3)	(1.2)	(0.7)	(1.1)	(100.0)				
150～	1	1	4	3	3	9	7	1	4	2	1	1				37		37	5.21	1.14
(%)	(2.7)	(2.7)	(10.8)	(8.1)	(8.1)	(24.3)	(18.9)	(2.7)	(10.8)	(5.4)	(2.7)	(2.7)				(100.0)				
151～	14	29	60	95	148	163	147	29	94	65	42	22	9	6	8	931	3	934	5.38	1.22
(%)	(1.5)	(3.1)	(6.4)	(10.2)	(15.9)	(17.5)	(15.8)	(3.1)	(10.1)	(7.0)	(4.5)	(2.4)	(1.0)	(0.6)	(0.9)	(100.0)				
180～	14	16	36	70	107	111	97	20	65	49	26	22	17	5	7	662	4	666	5.45	1.31
(%)	(2.1)	(2.4)	(5.4)	(10.6)	(16.2)	(16.8)	(14.7)	(3.0)	(9.8)	(7.4)	(3.9)	(3.3)	(2.6)	(0.8)	(1.1)	(100.0)				
210～	5	22	23	52	78	66	76	11	40	32	25	13	8	6	15	472		472	5.51	1.43
(%)	(1.1)	(4.7)	(4.9)	(11.0)	(16.5)	(14.0)	(16.1)	(2.3)	(8.5)	(6.8)	(5.3)	(2.8)	(1.7)	(1.3)	(3.2)	(100.0)				
240～	12	18	32	46	67	80	86	12	39	49	33	25	12	1	13	525	1	526	5.58	1.41
(%)	(2.3)	(3.4)	(6.1)	(8.8)	(12.8)	(15.2)	(16.4)	(2.3)	(7.4)	(9.3)	(6.3)	(4.8)	(2.3)	(0.2)	(2.5)	(100.0)				
300～	2	4	11	15	30	33	35	3	17	22	14	9	2	4	4	205		205	5.70	1.35
(%)	(1.0)	(2.0)	(5.4)	(7.3)	(14.6)	(16.1)	(17.1)	(1.5)	(8.3)	(10.7)	(6.8)	(4.4)	(1.0)	(2.0)	(2.0)	(100.0)				
350～	1	3	7	8	19	20	24	4	13	11	8	8	2	1	4	133	2	135	5.76	1.41
(%)	(0.8)	(2.3)	(5.3)	(6.0)	(14.3)	(15.0)	(18.0)	(3.0)	(9.8)	(8.3)	(6.0)	(6.0)	(1.5)	(0.8)	(3.0)	(100.0)				
400～	2	8	4	12	20	16	22	4	15	14	16	7	2	4	5	151	1	152	5.84	1.58
(%)	(1.3)	(5.3)	(2.6)	(7.9)	(13.2)	(10.6)	(14.6)	(2.6)	(9.9)	(9.3)	(10.6)	(4.6)	(1.3)	(2.6)	(3.3)	(100.0)				
500～	1	7	6	16	16	25	12	6	15	13	11	10	6	3	19	166		166	6.24	2.11
(%)	(0.6)	(4.2)	(3.6)	(9.6)	(9.6)	(15.1)	(7.2)	(3.6)	(9.0)	(7.8)	(6.6)	(6.0)	(3.6)	(1.8)	(11.4)	(100.0)				
合計	613	864	1,480	1,986	2,499	2,535	2,041	345	1,123	963	538	332	179	93	178	15,769	65	15,834	5.11	1.36
(%)	(3.9)	(5.5)	(9.4)	(12.6)	(15.8)	(16.1)	(12.9)	(2.2)	(7.1)	(6.1)	(3.4)	(2.1)	(1.1)	(0.6)	(1.1)	(100.0)				
記載なし	28	29	35	45	34	39	23	3	11	12	10	7	2	1	1	280	158	438	4.65	1.42
(%)	(10.0)	(10.4)	(12.5)	(16.1)	(12.1)	(13.9)	(8.2)	(1.1)	(3.9)	(4.3)	(3.6)	(2.5)	(0.7)	(0.4)	(0.4)	(100.0)				
総計	641	893	1,515	2,031	2,533	2,574	2,064	348	1,134	975	548	339	181	94	179	16,049	223	16,272	5.11	1.36
(%)	(4.0)	(5.6)	(9.4)	(12.7)	(15.8)	(16.0)	(12.9)	(2.2)	(7.1)	(6.1)	(3.4)	(2.1)	(1.1)	(0.6)	(1.1)	(100.0)				
平均	68.00	84.84	84.33	93.80	100.21	103.76	114.63	120.66	118.72	120.85	140.51	153.17	141.54	151.18	200.45	105.67	111.42	105.70		
標準偏差	74.84	90.48	89.51	97.28	91.25	94.72	122.27	129.42	106.08	109.52	140.08	185.50	150.68	148.13	226.57	109.34	81.01	109.24		

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査