

表 3463 透析前リン濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別(施設血液透析患者 透析回数週3回)

透析前ヘモグロビン 濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
5未満	18	9	21	36	38	42	31	6	16	19	9	10	2	1	2	260	1	261	5.09	1.41
(%)	(6.9)	(3.5)	(8.1)	(13.8)	(14.6)	(16.2)	(11.9)	(2.3)	(6.2)	(7.3)	(3.5)	(3.8)	(0.8)	(0.4)	(0.8)	(100.0)				
5～	31	24	28	35	51	53	39	4	26	29	19	5	7	2	7	360	2	362	5.10	1.64
(%)	(8.6)	(6.7)	(7.8)	(9.7)	(14.2)	(14.7)	(10.8)	(1.1)	(7.2)	(8.1)	(5.3)	(1.4)	(1.9)	(0.6)	(1.9)	(100.0)				
6～	137	78	98	102	112	104	93	16	45	43	31	18	13	14	18	922	20	942	4.74	1.80
(%)	(14.9)	(8.5)	(10.6)	(11.1)	(12.1)	(11.3)	(10.1)	(1.7)	(4.9)	(4.7)	(3.4)	(2.0)	(1.4)	(1.5)	(2.0)	(100.0)				
7～	444	308	418	477	468	392	334	47	193	145	89	77	51	27	50	3,520	58	3,578	4.70	1.67
(%)	(12.6)	(8.8)	(11.9)	(13.6)	(13.3)	(11.1)	(9.5)	(1.3)	(5.5)	(4.1)	(2.5)	(2.2)	(1.4)	(0.8)	(1.4)	(100.0)				
8～	1,109	939	1,335	1,612	1,648	1,461	1,169	213	626	535	358	213	152	116	139	11,625	101	11,726	4.80	1.55
(%)	(9.5)	(8.1)	(11.5)	(13.9)	(14.2)	(12.6)	(10.1)	(1.8)	(5.4)	(4.6)	(3.1)	(1.8)	(1.3)	(1.0)	(1.2)	(100.0)				
9～	2,123	2,345	3,730	5,171	5,540	5,328	4,276	713	2,360	2,054	1,229	803	453	296	455	36,876	162	37,038	5.01	1.44
(%)	(5.8)	(6.4)	(10.1)	(14.0)	(15.0)	(14.4)	(11.6)	(1.9)	(6.4)	(5.6)	(3.3)	(2.2)	(1.2)	(0.8)	(1.2)	(100.0)				
10～	2,503	3,594	6,618	9,946	11,941	12,007	9,912	1,683	5,583	4,581	2,778	1,562	916	556	868	75,048	169	75,217	5.16	1.35
(%)	(3.3)	(4.8)	(8.8)	(13.3)	(15.9)	(16.0)	(13.2)	(2.2)	(7.4)	(6.1)	(3.7)	(2.1)	(1.2)	(0.7)	(1.2)	(100.0)				
11～	1,704	2,475	4,698	7,374	9,384	9,735	8,700	1,443	4,910	4,431	2,618	1,603	878	559	860	61,372	107	61,479	5.31	1.38
(%)	(2.8)	(4.0)	(7.7)	(12.0)	(15.3)	(15.9)	(14.2)	(2.4)	(8.0)	(7.2)	(4.3)	(2.6)	(1.4)	(0.9)	(1.4)	(100.0)				
12～	582	797	1,491	2,475	3,046	3,415	3,161	574	1,936	1,776	1,191	683	490	282	421	22,320	50	22,370	5.46	1.44
(%)	(2.6)	(3.6)	(6.7)	(11.1)	(13.6)	(15.3)	(14.2)	(2.6)	(8.7)	(8.0)	(5.3)	(3.1)	(2.2)	(1.3)	(1.9)	(100.0)				
13～	143	217	296	517	702	726	783	137	502	510	343	221	127	99	196	5,519	18	5,537	5.67	1.59
(%)	(2.6)	(3.9)	(5.4)	(9.4)	(12.7)	(13.2)	(14.2)	(2.5)	(9.1)	(9.2)	(6.2)	(4.0)	(2.3)	(1.8)	(3.6)	(100.0)				
14～	58	68	101	160	230	246	240	44	173	144	127	103	74	50	109	1,927	6	1,933	5.85	1.76
(%)	(3.0)	(3.5)	(5.2)	(8.3)	(11.9)	(12.8)	(12.5)	(2.3)	(9.0)	(7.5)	(6.6)	(5.3)	(3.8)	(2.6)	(5.7)	(100.0)				
合計	8,852	10,854	18,834	27,905	33,160	33,509	28,738	4,880	16,370	14,267	8,792	5,298	3,163	2,002	3,125	219,749	694	220,443	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	109	126	184	287	308	300	274	48	135	110	68	61	29	21	31	2,091	6,437	8,528	5.11	1.48
(%)	(5.2)	(6.0)	(8.8)	(13.7)	(14.7)	(14.3)	(13.1)	(2.3)	(6.5)	(5.3)	(3.3)	(2.9)	(1.4)	(1.0)	(1.5)	(100.0)				
総計	8,961	10,980	19,018	28,192	33,468	33,809	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
平均	10.12	10.39	10.49	10.59	10.67	10.73	10.80	10.82	10.84	10.88	10.92	10.93	10.94	10.94	11.01	10.69	9.95	10.69		
標準偏差	1.46	1.33	1.25	1.23	1.22	1.21	1.22	1.23	1.24	1.27	1.32	1.38	1.43	1.45	1.53	1.28	1.65	1.28		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査