

表 3465 透析前リン濃度(mg/dL) HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

HDL-C濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	133	56	67	80	86	69	61	6	37	30	12	14	7	3	6	667	4	671	4.41	1.77
(%)	(19.9)	(8.4)	(10.0)	(12.0)	(12.9)	(10.3)	(9.1)	(0.9)	(5.5)	(4.5)	(1.8)	(2.1)	(1.0)	(0.4)	(0.9)	(100.0)				
20～	719	786	1,206	1,728	1,830	1,887	1,582	248	876	755	477	282	158	123	166	12,823	23	12,846	5.07	1.47
(%)	(5.6)	(6.1)	(9.4)	(13.5)	(14.3)	(14.7)	(12.3)	(1.9)	(6.8)	(5.9)	(3.7)	(2.2)	(1.2)	(1.0)	(1.3)	(100.0)				
30～	1,604	1,954	3,320	4,879	5,685	5,532	4,711	808	2,582	2,220	1,408	825	514	316	487	36,845	43	36,888	5.14	1.42
(%)	(4.4)	(5.3)	(9.0)	(13.2)	(15.4)	(15.0)	(12.8)	(2.2)	(7.0)	(6.0)	(3.8)	(2.2)	(1.4)	(0.9)	(1.3)	(100.0)				
40～	1,419	1,956	3,578	5,315	6,314	6,410	5,363	895	3,039	2,572	1,565	922	513	321	504	40,686	43	40,729	5.18	1.38
(%)	(3.5)	(4.8)	(8.8)	(13.1)	(15.5)	(15.8)	(13.2)	(2.2)	(7.5)	(6.3)	(3.8)	(2.3)	(1.3)	(0.8)	(1.2)	(100.0)				
50～	957	1,280	2,418	3,541	4,445	4,614	4,084	688	2,232	1,892	1,171	679	414	259	382	29,056	24	29,080	5.24	1.38
(%)	(3.3)	(4.4)	(8.3)	(12.2)	(15.3)	(15.9)	(14.1)	(2.4)	(7.7)	(6.5)	(4.0)	(2.3)	(1.4)	(0.9)	(1.3)	(100.0)				
60～	471	704	1,320	2,054	2,558	2,721	2,270	385	1,281	1,162	739	421	255	162	269	16,772	20	16,792	5.30	1.39
(%)	(2.8)	(4.2)	(7.9)	(12.2)	(15.3)	(16.2)	(13.5)	(2.3)	(7.6)	(6.9)	(4.4)	(2.5)	(1.5)	(1.0)	(1.6)	(100.0)				
70～	217	332	611	1,041	1,187	1,267	1,119	198	679	594	372	229	131	79	134	8,190	11	8,201	5.34	1.40
(%)	(2.6)	(4.1)	(7.5)	(12.7)	(14.5)	(15.5)	(13.7)	(2.4)	(8.3)	(7.3)	(4.5)	(2.8)	(1.6)	(1.0)	(1.6)	(100.0)				
80～	94	136	288	468	589	585	511	108	311	292	160	107	68	42	71	3,830	4	3,834	5.37	1.43
(%)	(2.5)	(3.6)	(7.5)	(12.2)	(15.4)	(15.3)	(13.3)	(2.8)	(8.1)	(7.6)	(4.2)	(2.8)	(1.8)	(1.1)	(1.9)	(100.0)				
90～	44	69	115	216	278	239	235	42	148	119	81	59	32	17	37	1,731	2	1,733	5.41	1.49
(%)	(2.5)	(4.0)	(6.6)	(12.5)	(16.1)	(13.8)	(13.6)	(2.4)	(8.5)	(6.9)	(4.7)	(3.4)	(1.8)	(1.0)	(2.1)	(100.0)				
100～	43	65	98	165	208	217	199	34	130	109	61	39	31	26	31	1,456	2	1,458	5.41	1.45
(%)	(3.0)	(4.5)	(6.7)	(11.3)	(14.3)	(14.9)	(13.7)	(2.3)	(8.9)	(7.5)	(4.2)	(2.7)	(2.1)	(1.8)	(2.1)	(100.0)				
合計	5,701	7,338	13,021	19,487	23,180	23,541	20,135	3,412	11,315	9,745	6,046	3,577	2,123	1,348	2,087	152,056	176	152,232	5.20	1.41
(%)	(3.7)	(4.8)	(8.6)	(12.8)	(15.2)	(15.5)	(13.2)	(2.2)	(7.4)	(6.4)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	3,260	3,642	5,997	8,705	10,288	10,268	8,877	1,516	5,190	4,632	2,814	1,782	1,069	675	1,069	69,784	6,955	76,739	5.19	1.47
(%)	(4.7)	(5.2)	(8.6)	(12.5)	(14.7)	(14.7)	(12.7)	(2.2)	(7.4)	(6.6)	(4.0)	(2.6)	(1.5)	(1.0)	(1.5)	(100.0)				
総計	8,961	10,980	19,018	28,192	33,468	33,809	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
平均	44.50	46.32	47.13	47.71	48.24	48.55	48.73	49.19	49.19	49.41	49.17	49.53	49.84	49.80	50.32	48.26	46.69	48.25		
標準偏差	16.32	16.31	16.10	16.25	16.32	16.32	16.30	16.47	16.98	17.09	16.69	17.29	17.70	18.51	18.57	16.54	18.50	16.54		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査