

表 3466 透析前リン濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

Non-HDL-C濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
50未満	198	178	285	322	396	398	344	49	152	142	88	58	34	32	34	2,710	6	2,716	4.99	1.50
(%)	(7.3)	(6.6)	(10.5)	(11.9)	(14.6)	(14.7)	(12.7)	(1.8)	(5.6)	(5.2)	(3.2)	(2.1)	(1.3)	(1.2)	(1.3)	(100.0)				
50～	699	673	1,243	1,734	1,998	1,934	1,607	265	874	721	456	262	182	93	148	12,889	17	12,906	5.06	1.42
(%)	(5.4)	(5.2)	(9.6)	(13.5)	(15.5)	(15.0)	(12.5)	(2.1)	(6.8)	(5.6)	(3.5)	(2.0)	(1.4)	(0.7)	(1.1)	(100.0)				
70～	1,247	1,556	2,607	3,849	4,440	4,379	3,627	587	1,977	1,687	1,093	577	369	192	313	28,500	32	28,532	5.10	1.38
(%)	(4.4)	(5.5)	(9.1)	(13.5)	(15.6)	(15.4)	(12.7)	(2.1)	(6.9)	(5.9)	(3.8)	(2.0)	(1.3)	(0.7)	(1.1)	(100.0)				
90～	1,219	1,667	2,941	4,441	5,142	5,217	4,440	780	2,507	2,038	1,291	770	394	284	422	33,553	26	33,579	5.17	1.39
(%)	(3.6)	(5.0)	(8.8)	(13.2)	(15.3)	(15.5)	(13.2)	(2.3)	(7.5)	(6.1)	(3.8)	(2.3)	(1.2)	(0.8)	(1.3)	(100.0)				
110～	850	1,227	2,242	3,295	4,103	4,170	3,627	634	2,089	1,740	1,035	604	391	259	388	26,654	25	26,679	5.24	1.39
(%)	(3.2)	(4.6)	(8.4)	(12.4)	(15.4)	(15.6)	(13.6)	(2.4)	(7.8)	(6.5)	(3.9)	(2.3)	(1.5)	(1.0)	(1.5)	(100.0)				
130～	475	664	1,220	1,943	2,423	2,561	2,172	368	1,295	1,143	703	471	258	179	243	16,118	15	16,133	5.32	1.41
(%)	(2.9)	(4.1)	(7.6)	(12.1)	(15.0)	(15.9)	(13.5)	(2.3)	(8.0)	(7.1)	(4.4)	(2.9)	(1.6)	(1.1)	(1.5)	(100.0)				
150～	221	294	601	959	1,171	1,237	1,079	189	662	610	401	223	134	92	163	8,036	7	8,043	5.39	1.45
(%)	(2.8)	(3.7)	(7.5)	(11.9)	(14.6)	(15.4)	(13.4)	(2.4)	(8.2)	(7.6)	(5.0)	(2.8)	(1.7)	(1.1)	(2.0)	(100.0)				
170～	107	134	254	392	519	521	475	92	267	292	205	106	81	44	72	3,561	6	3,567	5.44	1.49
(%)	(3.0)	(3.8)	(7.1)	(11.0)	(14.6)	(14.6)	(13.3)	(2.6)	(7.5)	(8.2)	(5.8)	(3.0)	(2.3)	(1.2)	(2.0)	(100.0)				
190～	45	51	107	150	186	182	182	27	108	106	71	44	29	10	38	1,336	2	1,338	5.43	1.55
(%)	(3.4)	(3.8)	(8.0)	(11.2)	(13.9)	(13.6)	(13.6)	(2.0)	(8.1)	(7.9)	(5.3)	(3.3)	(2.2)	(0.7)	(2.8)	(100.0)				
210～	27	38	55	73	102	129	113	10	65	69	48	36	20	13	32	830	2	832	5.60	1.69
(%)	(3.3)	(4.6)	(6.6)	(8.8)	(12.3)	(15.5)	(13.6)	(1.2)	(7.8)	(8.3)	(5.8)	(4.3)	(2.4)	(1.6)	(3.9)	(100.0)				
合計	5,088	6,482	11,555	17,158	20,480	20,728	17,666	3,001	9,996	8,548	5,391	3,151	1,892	1,198	1,853	134,187	138	134,325	5.20	1.41
(%)	(3.8)	(4.8)	(8.6)	(12.8)	(15.3)	(15.4)	(13.2)	(2.2)	(7.4)	(6.4)	(4.0)	(2.3)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	3,873	4,498	7,463	11,034	12,988	13,081	11,346	1,927	6,509	5,829	3,469	2,208	1,300	825	1,303	87,653	6,993	94,646	5.19	1.45
(%)	(4.4)	(5.1)	(8.5)	(12.6)	(14.8)	(14.9)	(12.9)	(2.2)	(7.4)	(6.7)	(4.0)	(2.5)	(1.5)	(0.9)	(1.5)	(100.0)				
総計	8,961	10,980	19,018	28,192	33,468	33,809	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
平均	99.73	102.47	103.71	104.75	105.77	106.57	107.26	107.77	108.43	109.87	110.21	111.21	110.71	111.89	113.72	106.44	104.46	106.43		
標準偏差	33.71	32.30	32.86	32.17	32.68	32.85	33.14	32.27	33.25	34.28	34.98	35.68	36.52	34.87	38.22	33.31	38.86	33.32		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査