

表 3467 透析前リン濃度(mg/dL) 透析前収縮期血圧(mmHg)別（施設血液透析患者 透析回数週3回）

透析前収縮期血圧	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
60未満	3	4	5	4	2	4	3			1	3	2		2	1	34	1	35	5.05	1.90
(%)	(8.8)	(11.8)	(14.7)	(11.8)	(5.9)	(11.8)	(8.8)			(2.9)	(8.8)	(5.9)		(5.9)	(2.9)	(100.0)				
60～	19	15	21	36	41	37	26	8	14	15	10	7	6	3	8	266	5	271	5.13	1.71
(%)	(7.1)	(5.6)	(7.9)	(13.5)	(15.4)	(13.9)	(9.8)	(3.0)	(5.3)	(5.6)	(3.8)	(2.6)	(2.3)	(1.1)	(3.0)	(100.0)				
80～	187	148	215	293	303	340	275	50	146	139	87	51	34	27	38	2,333	44	2,377	5.03	1.59
(%)	(8.0)	(6.3)	(9.2)	(12.6)	(13.0)	(14.6)	(11.8)	(2.1)	(6.3)	(6.0)	(3.7)	(2.2)	(1.5)	(1.2)	(1.6)	(100.0)				
100～	764	769	1,205	1,704	1,936	1,820	1,598	276	941	741	507	269	178	98	155	12,961	194	13,155	5.06	1.46
(%)	(5.9)	(5.9)	(9.3)	(13.1)	(14.9)	(14.0)	(12.3)	(2.1)	(7.3)	(5.7)	(3.9)	(2.1)	(1.4)	(0.8)	(1.2)	(100.0)				
120～	1,758	2,207	3,708	5,368	6,306	6,353	5,323	930	2,996	2,502	1,467	916	539	321	476	41,170	406	41,576	5.13	1.39
(%)	(4.3)	(5.4)	(9.0)	(13.0)	(15.3)	(15.4)	(12.9)	(2.3)	(7.3)	(6.1)	(3.6)	(2.2)	(1.3)	(0.8)	(1.2)	(100.0)				
140～	2,414	3,134	5,517	8,151	9,812	10,001	8,524	1,397	4,700	4,030	2,417	1,482	830	521	795	63,725	478	64,203	5.18	1.38
(%)	(3.8)	(4.9)	(8.7)	(12.8)	(15.4)	(15.7)	(13.4)	(2.2)	(7.4)	(6.3)	(3.8)	(2.3)	(1.3)	(0.8)	(1.2)	(100.0)				
160～	1,604	2,046	3,910	5,902	7,017	7,141	6,185	1,050	3,520	3,197	1,978	1,144	725	411	693	46,523	320	46,843	5.25	1.42
(%)	(3.4)	(4.4)	(8.4)	(12.7)	(15.1)	(15.3)	(13.3)	(2.3)	(7.6)	(6.9)	(4.3)	(2.5)	(1.6)	(0.9)	(1.5)	(100.0)				
180～	624	844	1,378	2,158	2,556	2,647	2,343	413	1,412	1,308	817	513	293	219	361	17,886	116	18,002	5.33	1.49
(%)	(3.5)	(4.7)	(7.7)	(12.1)	(14.3)	(14.8)	(13.1)	(2.3)	(7.9)	(7.3)	(4.6)	(2.9)	(1.6)	(1.2)	(2.0)	(100.0)				
200～	143	196	319	559	595	655	622	104	341	313	239	122	97	63	115	4,483	32	4,515	5.42	1.54
(%)	(3.2)	(4.4)	(7.1)	(12.5)	(13.3)	(14.6)	(13.9)	(2.3)	(7.6)	(7.0)	(5.3)	(2.7)	(2.2)	(1.4)	(2.6)	(100.0)				
220～	28	32	65	91	125	132	107	14	67	68	51	25	25	19	33	882	12	894	5.54	1.64
(%)	(3.2)	(3.6)	(7.4)	(10.3)	(14.2)	(15.0)	(12.1)	(1.6)	(7.6)	(7.7)	(5.8)	(2.8)	(2.8)	(2.2)	(3.7)	(100.0)				
合計	7,544	9,395	16,343	24,266	28,693	29,130	25,006	4,242	14,137	12,314	7,576	4,531	2,727	1,684	2,675	190,263	1,608	191,871	5.20	1.42
(%)	(4.0)	(4.9)	(8.6)	(12.8)	(15.1)	(15.3)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	1,417	1,585	2,675	3,926	4,775	4,679	4,006	686	2,368	2,063	1,284	828	465	339	481	31,577	5,523	37,100	5.21	1.46
(%)	(4.5)	(5.0)	(8.5)	(12.4)	(15.1)	(14.8)	(12.7)	(2.2)	(7.5)	(6.5)	(4.1)	(2.6)	(1.5)	(1.1)	(1.5)	(100.0)				
総計	8,961	10,980	19,018	28,192	33,468	33,809	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
平均	147.70	149.54	150.44	151.20	151.38	151.70	152.16	151.98	152.35	153.27	153.60	153.60	154.41	155.23	156.30	151.69	146.21	151.64		
標準偏差	25.23	24.28	23.76	23.75	23.45	23.55	23.65	23.95	23.93	24.11	24.86	24.39	25.71	26.27	26.99	23.99	26.22	24.02		

数値下のかつこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査