

表 3469 透析前リン濃度(mg/dL) 透析前脈拍(拍/分)別（施設血液透析患者 透析回数週3回）

透析前脈拍	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
40未満	8	15	16	17	16	23	12	4	9	15	10	4	1	3	2	155		155	5.13	1.60
(%)	(5.2)	(9.7)	(10.3)	(11.0)	(10.3)	(14.8)	(7.7)	(2.6)	(5.8)	(9.7)	(6.5)	(2.6)	(0.6)	(1.9)	(1.3)	(100.0)				
40～	94	110	173	264	294	310	223	39	119	119	65	32	18	18	29	1,907	22	1,929	5.06	1.41
(%)	(4.9)	(5.8)	(9.1)	(13.8)	(15.4)	(16.3)	(11.7)	(2.0)	(6.2)	(6.2)	(3.4)	(1.7)	(0.9)	(0.9)	(1.5)	(100.0)				
50～	679	814	1,480	2,132	2,464	2,487	1,981	316	1,130	922	527	327	163	110	109	15,641	132	15,773	5.07	1.33
(%)	(4.3)	(5.2)	(9.5)	(13.6)	(15.8)	(15.9)	(12.7)	(2.0)	(7.2)	(5.9)	(3.4)	(2.1)	(1.0)	(0.7)	(0.7)	(100.0)				
60～	1,827	2,477	4,311	6,279	7,617	7,600	6,428	1,071	3,583	2,998	1,764	1,010	589	332	478	48,364	377	48,741	5.14	1.35
(%)	(3.8)	(5.1)	(8.9)	(13.0)	(15.7)	(15.7)	(13.3)	(2.2)	(7.4)	(6.2)	(3.6)	(2.1)	(1.2)	(0.7)	(1.0)	(100.0)				
70～	2,142	2,843	4,877	7,385	8,866	8,812	7,750	1,316	4,270	3,737	2,271	1,406	776	506	765	57,722	462	58,184	5.20	1.40
(%)	(3.7)	(4.9)	(8.4)	(12.8)	(15.4)	(15.3)	(13.4)	(2.3)	(7.4)	(6.5)	(3.9)	(2.4)	(1.3)	(0.9)	(1.3)	(100.0)				
80～	1,462	1,708	3,020	4,541	5,252	5,492	4,709	819	2,787	2,423	1,613	900	632	368	658	36,384	339	36,723	5.26	1.48
(%)	(4.0)	(4.7)	(8.3)	(12.5)	(14.4)	(15.1)	(12.9)	(2.3)	(7.7)	(6.7)	(4.4)	(2.5)	(1.7)	(1.0)	(1.8)	(100.0)				
90～	661	737	1,320	1,920	2,140	2,297	2,037	344	1,230	1,120	687	468	304	183	345	15,793	125	15,918	5.31	1.53
(%)	(4.2)	(4.7)	(8.4)	(12.2)	(13.6)	(14.5)	(12.9)	(2.2)	(7.8)	(7.1)	(4.4)	(3.0)	(1.9)	(1.2)	(2.2)	(100.0)				
100～	245	241	395	573	673	657	566	128	339	357	244	144	93	68	130	4,853	51	4,904	5.32	1.62
(%)	(5.0)	(5.0)	(8.1)	(11.8)	(13.9)	(13.5)	(11.7)	(2.6)	(7.0)	(7.4)	(5.0)	(3.0)	(1.9)	(1.4)	(2.7)	(100.0)				
110～	70	57	98	140	159	165	144	21	80	51	57	30	21	16	33	1,142	15	1,157	5.22	1.62
(%)	(6.1)	(5.0)	(8.6)	(12.3)	(13.9)	(14.4)	(12.6)	(1.8)	(7.0)	(4.5)	(5.0)	(2.6)	(1.8)	(1.4)	(2.9)	(100.0)				
120～	37	20	33	65	64	76	61	10	30	34	21	14	12	6	13	496	2	498	5.25	1.62
(%)	(7.5)	(4.0)	(6.7)	(13.1)	(12.9)	(15.3)	(12.3)	(2.0)	(6.0)	(6.9)	(4.2)	(2.8)	(2.4)	(1.2)	(2.6)	(100.0)				
合計	7,225	9,022	15,723	23,316	27,545	27,919	23,911	4,068	13,577	11,776	7,259	4,335	2,609	1,610	2,562	182,457	1,525	183,982	5.19	1.42
(%)	(4.0)	(4.9)	(8.6)	(12.8)	(15.1)	(15.3)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
記載なし	1,736	1,958	3,295	4,876	5,923	5,890	5,101	860	2,928	2,601	1,601	1,024	583	413	594	39,383	5,606	44,989	5.21	1.46
(%)	(4.4)	(5.0)	(8.4)	(12.4)	(15.0)	(15.0)	(13.0)	(2.2)	(7.4)	(6.6)	(4.1)	(2.6)	(1.5)	(1.0)	(1.5)	(100.0)				
総計	8,961	10,980	19,018	28,192	33,468	33,809	29,012	4,928	16,505	14,377	8,860	5,359	3,192	2,023	3,156	221,840	7,131	228,971	5.20	1.43
(%)	(4.0)	(4.9)	(8.6)	(12.7)	(15.1)	(15.2)	(13.1)	(2.2)	(7.4)	(6.5)	(4.0)	(2.4)	(1.4)	(0.9)	(1.4)	(100.0)				
平均	74.98	74.00	74.02	74.13	73.94	74.12	74.37	74.68	74.69	75.01	75.66	75.94	76.90	76.88	78.73	74.49	74.99	74.50		
標準偏差	13.78	12.77	12.79	12.76	12.62	12.71	12.49	12.56	12.59	12.89	12.97	12.86	13.22	13.65	13.56	12.80	13.12	12.80		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査