

表 3473 透析前リン濃度(mg/dL) 原疾患別 (血液透析濾過患者 透析回数週3回)

原疾患	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	373	540	985	1,718	2,215	2,458	2,068	393	1,220	1,060	656	416	247	151	235	14,735	341	15,076	5.37	1.38
(%)	(2.5)	(3.7)	(6.7)	(11.7)	(15.0)	(16.7)	(14.0)	(2.7)	(8.3)	(7.2)	(4.5)	(2.8)	(1.7)	(1.0)	(1.6)	(100.0)				
慢性腎盂腎炎	13	11	30	54	48	62	60	10	44	35	16	5	4	9	10	411	11	422	5.42	1.44
(%)	(3.2)	(2.7)	(7.3)	(13.1)	(11.7)	(15.1)	(14.6)	(2.4)	(10.7)	(8.5)	(3.9)	(1.2)	(1.0)	(2.2)	(2.4)	(100.0)				
急速進行性糸球体腎炎	7	12	21	36	51	49	30	11	14	13	14	7	5	4	7	281	3	284	5.31	1.47
(%)	(2.5)	(4.3)	(7.5)	(12.8)	(18.1)	(17.4)	(10.7)	(3.9)	(5.0)	(4.6)	(5.0)	(2.5)	(1.8)	(1.4)	(2.5)	(100.0)				
妊娠腎/妊娠中毒症	5	14	25	44	57	58	57	11	28	22	17	4	2	4	4	352	7	359	5.30	1.29
(%)	(1.4)	(4.0)	(7.1)	(12.5)	(16.2)	(16.5)	(16.2)	(3.1)	(8.0)	(6.3)	(4.8)	(1.1)	(0.6)	(1.1)	(1.1)	(100.0)				
その他の分類不能の腎炎	6	3	13	16	27	32	19	5	23	10	9	5	8	4	10	190	5	195	5.71	1.70
(%)	(3.2)	(1.6)	(6.8)	(8.4)	(14.2)	(16.8)	(10.0)	(2.6)	(12.1)	(5.3)	(4.7)	(2.6)	(4.2)	(2.1)	(5.3)	(100.0)				
多発性囊胞腎	30	45	81	161	199	246	195	40	136	106	72	47	22	13	20	1,413	51	1,464	5.45	1.33
(%)	(2.1)	(3.2)	(5.7)	(11.4)	(14.1)	(17.4)	(13.8)	(2.8)	(9.6)	(7.5)	(5.1)	(3.3)	(1.6)	(0.9)	(1.4)	(100.0)				
腎硬化症	114	136	233	354	423	414	373	71	226	186	111	75	39	24	45	2,824	62	2,886	5.23	1.46
(%)	(4.0)	(4.8)	(8.3)	(12.5)	(15.0)	(14.7)	(13.2)	(2.5)	(8.0)	(6.6)	(3.9)	(2.7)	(1.4)	(0.8)	(1.6)	(100.0)				
悪性高血圧	5	7	17	31	34	58	36	7	26	26	19	9	9	7	11	302	10	312	5.70	1.55
(%)	(1.7)	(2.3)	(5.6)	(10.3)	(11.3)	(19.2)	(11.9)	(2.3)	(8.6)	(8.6)	(6.3)	(3.0)	(3.0)	(2.3)	(3.6)	(100.0)				
糖尿病性腎症	411	565	1,056	1,523	1,886	1,870	1,678	266	1,046	938	608	373	210	145	242	12,817	356	13,173	5.33	1.46
(%)	(3.2)	(4.4)	(8.2)	(11.9)	(14.7)	(14.6)	(13.1)	(2.1)	(8.2)	(7.3)	(4.7)	(2.9)	(1.6)	(1.1)	(1.9)	(100.0)				
SLE腎炎	12	7	19	33	38	73	59	10	21	29	16	7	14	1	13	352	11	363	5.59	1.58
(%)	(3.4)	(2.0)	(5.4)	(9.4)	(10.8)	(20.7)	(16.8)	(2.8)	(6.0)	(8.2)	(4.5)	(2.0)	(4.0)	(0.3)	(3.7)	(100.0)				
アミロイド腎	1	3	8	6	8	11	5	4	4	3	2	2	2			59		59	5.17	1.29
(%)	(1.7)	(5.1)	(13.6)	(10.2)	(13.6)	(18.6)	(8.5)	(6.8)	(6.8)	(5.1)	(3.4)	(3.4)	(3.4)			(100.0)				
痛風腎	3	7	14	16	19	22	16	4	12	9	7	9	2	3	3	146	8	154	5.44	1.50
(%)	(2.1)	(4.8)	(9.6)	(11.0)	(13.0)	(15.1)	(11.0)	(2.7)	(8.2)	(6.2)	(4.8)	(6.2)	(1.4)	(2.1)	(2.1)	(100.0)				
先天性代謝異常 による腎不全		2	1	6	3	7	6	2	3	4	4		1		2	41	3	44	5.71	1.49
(%)		(4.9)	(2.4)	(14.6)	(7.3)	(17.1)	(14.6)	(4.9)	(7.3)	(9.8)	(9.8)		(2.4)		(4.9)	(100.0)				
腎・尿路結核			2	5	2	6	2	1				1	1	1		21	1	22	5.21	1.22
(%)			(9.5)	(23.8)	(9.5)	(28.6)	(9.5)	(4.8)				(4.8)	(4.8)	(4.8)		(100.0)				
腎・尿路結石	2	4	9	4	7	12	13	1	5	3	5		1		2	68	1	69	5.30	1.60
(%)	(2.9)	(5.9)	(13.2)	(5.9)	(10.3)	(17.6)	(19.1)	(1.5)	(7.4)	(4.4)	(7.4)		(1.5)		(2.9)	(100.0)				
腎・尿路腫瘍	3	3	7	9	16	11	11	3	2	7	3	4	4	1	1	85		85	5.38	1.53
(%)	(3.5)	(3.5)	(8.2)	(10.6)	(18.8)	(12.9)	(12.9)	(3.5)	(2.4)	(8.2)	(3.5)	(4.7)	(4.7)	(1.2)	(1.2)	(100.0)				
閉塞性尿路障害	1	3	4	6	10	13	7	1	7	10	1	4	1	2	1	71	3	74	5.58	1.40
(%)	(1.4)	(4.2)	(5.6)	(8.5)	(14.1)	(18.3)	(9.9)	(1.4)	(9.9)	(14.1)	(1.4)	(5.6)	(1.4)	(2.8)	(1.4)	(100.0)				
骨髄腫	2	1	3	3	5	2	2					1				19		19	4.40	1.21
(%)	(10.5)	(5.3)	(15.8)	(15.8)	(26.3)	(10.5)	(10.5)					(5.3)				(100.0)				
腎形成不全	1	3	7	11	8	13	23	1	9	12	4	1	4	1	1	99	2	101	5.56	1.32
(%)	(1.0)	(3.0)	(7.1)	(11.1)	(8.1)	(13.1)	(23.2)	(1.0)	(9.1)	(12.1)	(4.0)	(1.0)	(4.0)	(1.0)	(1.0)	(100.0)				
不明	114	162	235	378	477	525	457	82	304	239	144	79	49	39	54	3,338	142	3,480	5.32	1.43
(%)	(3.4)	(4.9)	(7.0)	(11.3)	(14.3)	(15.7)	(13.7)	(2.5)	(9.1)	(7.2)	(4.3)	(2.4)	(1.5)	(1.2)	(1.6)	(100.0)				
再導入	6	10	24	48	59	55	63	6	38	39	20	15	9	8	7	407	9	416	5.55	1.37
(%)	(1.5)	(2.5)	(5.9)	(11.8)	(14.5)	(13.5)	(15.5)	(1.5)	(9.3)	(9.6)	(4.9)	(3.7)	(2.2)	(2.0)	(1.7)	(100.0)				
その他	24	36	55	103	120	137	119	21	76	55	46	40	17	7	14	870	19	889	5.42	1.46
(%)	(2.8)	(4.1)	(6.3)	(11.8)	(13.8)	(15.7)	(13.7)	(2.4)	(9.7)	(6.3)	(5.3)	(4.6)	(2.0)	(0.8)	(1.6)	(100.0)				
合計	1,133	1,574	2,849	4,565	5,712	6,134	5,299	950	3,244	2,806	1,775	1,104	651	423	682	38,901	1,045	39,946	5.36	1.43
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.8)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.8)	(1.7)	(1.1)	(1.8)	(100.0)				
記載なし																				
総計	1,133	1,574	2,849	4,565	5,712	6,134	5,299	950	3,244	2,806	1,775	1,104	651	423	682	38,901	1,045	39,946	5.36	1.43
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.8)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.8)	(1.7)	(1.1)	(1.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査