

表 3474 透析前リン濃度(mg/dL) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
国公立大学	8		6	2	3	5	6	4	1		1	1			1	38	4	42	4.67	1.80
(%)	(21.1)		(15.8)	(5.3)	(7.9)	(13.2)	(15.8)	(10.5)	(2.6)		(2.6)	(2.6)			(2.6)	(100.0)				
私立大学	5	4	11	18	38	33	19	7	12	8	4	3		2	1	165	8	173	5.16	1.22
(%)	(3.0)	(2.4)	(6.7)	(10.9)	(23.0)	(20.0)	(11.5)	(4.2)	(7.3)	(4.8)	(2.4)	(1.8)		(1.2)	(0.6)	(100.0)				
国立	6	4	3	4	5	7	4		1	1		1				36		36	4.39	1.40
(%)	(16.7)	(11.1)	(8.3)	(11.1)	(13.9)	(19.4)	(11.1)		(2.8)	(2.8)		(2.8)				(100.0)				
県市町村立	60	54	128	203	254	243	247	37	122	120	81	57	28	21	41	1,696	77	1,773	5.38	1.50
(%)	(3.5)	(3.2)	(7.5)	(12.0)	(15.0)	(14.3)	(14.6)	(2.2)	(7.2)	(7.1)	(4.8)	(3.4)	(1.7)	(1.2)	(2.4)	(100.0)				
社会保険	8	14	20	36	45	58	46	13	20	23	13	4	3	2	5	310	9	319	5.29	1.27
(%)	(2.6)	(4.5)	(6.5)	(11.6)	(14.5)	(18.7)	(14.8)	(4.2)	(6.5)	(7.4)	(4.2)	(1.3)	(1.0)	(0.6)	(1.6)	(100.0)				
厚生連	31	29	74	124	133	151	128	30	81	79	51	24	13	7	18	973	40	1,013	5.36	1.42
(%)	(3.2)	(3.0)	(7.6)	(12.7)	(13.7)	(15.5)	(13.2)	(3.1)	(8.3)	(8.1)	(5.2)	(2.5)	(1.3)	(0.7)	(1.8)	(100.0)				
その他公的	25	27	64	85	90	96	85	13	46	39	31	23	11	8	13	656	38	694	5.29	1.50
(%)	(3.8)	(4.1)	(9.8)	(13.0)	(13.7)	(14.6)	(13.0)	(2.0)	(7.0)	(5.9)	(4.7)	(3.5)	(1.7)	(1.2)	(2.0)	(100.0)				
私立総合	32	47	80	137	160	162	140	29	90	66	53	33	15	10	26	1,080	13	1,093	5.35	1.50
(%)	(3.0)	(4.4)	(7.4)	(12.7)	(14.8)	(15.0)	(13.0)	(2.7)	(8.3)	(6.1)	(4.9)	(3.1)	(1.4)	(0.9)	(2.4)	(100.0)				
私立	391	426	697	1,144	1,363	1,434	1,221	250	800	740	441	256	161	102	158	9,584	431	10,015	5.31	1.46
(%)	(4.1)	(4.4)	(7.3)	(11.9)	(14.2)	(15.0)	(12.7)	(2.6)	(8.3)	(7.7)	(4.6)	(2.7)	(1.7)	(1.1)	(1.6)	(100.0)				
私立診療所	567	969	1,766	2,812	3,621	3,945	3,403	567	2,071	1,730	1,100	702	420	271	419	24,363	425	24,788	5.38	1.40
(%)	(2.3)	(4.0)	(7.2)	(11.5)	(14.9)	(16.2)	(14.0)	(2.3)	(8.5)	(7.1)	(4.5)	(2.9)	(1.7)	(1.1)	(1.7)	(100.0)				
総計	1,133	1,574	2,849	4,565	5,712	6,134	5,299	950	3,244	2,806	1,775	1,104	651	423	682	38,901	1,045	39,946	5.36	1.43
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.8)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.8)	(1.7)	(1.1)	(1.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2014年調査