

表 3477 透析前リン濃度(mg/dL) PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	9	18	32	26	40	33	30	8	14	16	11	6	3	1	2	249		249	5.06	1.37
(%)	(3.6)	(7.2)	(12.9)	(10.4)	(16.1)	(13.3)	(12.0)	(3.2)	(5.6)	(6.4)	(4.4)	(2.4)	(1.2)	(0.4)	(0.8)	(100.0)				
20～	7	24	29	32	56	64	25	8	18	16	12	4	4	1	1	301		301	5.02	1.23
(%)	(2.3)	(8.0)	(9.6)	(10.6)	(18.6)	(21.3)	(8.3)	(2.7)	(6.0)	(5.3)	(4.0)	(1.3)	(1.3)	(0.3)	(0.3)	(100.0)				
35～	8	10	29	40	54	45	34	1	19	12	9	7		6		274		274	5.06	1.24
(%)	(2.9)	(3.6)	(10.6)	(14.6)	(19.7)	(16.4)	(12.4)	(0.4)	(6.9)	(4.4)	(3.3)	(2.6)		(2.2)		(100.0)				
50～	11	24	34	68	85	78	52	12	26	26	7	7	5	3	5	443	1	444	5.06	1.25
(%)	(2.5)	(5.4)	(7.7)	(15.3)	(19.2)	(17.6)	(11.7)	(2.7)	(5.9)	(5.9)	(1.6)	(1.6)	(1.1)	(0.7)	(1.1)	(100.0)				
75～	4	11	36	62	71	56	55	14	24	22	16	5	5	2	7	390	1	391	5.23	1.28
(%)	(1.0)	(2.8)	(9.2)	(15.9)	(18.2)	(14.4)	(14.1)	(3.6)	(6.2)	(5.6)	(4.1)	(1.3)	(1.3)	(0.5)	(1.8)	(100.0)				
100～	8	11	20	34	58	47	44	13	33	25	16	13	4	3	1	330	1	331	5.38	1.26
(%)	(2.4)	(3.3)	(6.1)	(10.3)	(17.6)	(14.2)	(13.3)	(3.9)	(10.0)	(7.6)	(4.8)	(3.9)	(1.2)	(0.9)	(0.3)	(100.0)				
125～	2	13	10	35	31	43	28	4	25	18	12	6	3		3	233	1	234	5.36	1.28
(%)	(0.9)	(5.6)	(4.3)	(15.0)	(13.3)	(18.5)	(12.0)	(1.7)	(10.7)	(7.7)	(5.2)	(2.6)	(1.3)		(1.3)	(100.0)				
150～					1		2						1			4		4	5.97	1.57
(%)					(25.0)		(50.0)						(25.0)			(100.0)				
151～	4	5	16	31	27	35	34	6	19	21	3	9		4	1	215		215	5.35	1.31
(%)	(1.9)	(2.3)	(7.4)	(14.4)	(12.6)	(16.3)	(15.8)	(2.8)	(8.8)	(9.8)	(1.4)	(4.2)		(1.9)	(0.5)	(100.0)				
180～	2	1	9	11	30	18	23	2	14	8	6	6	5		2	137		137	5.52	1.30
(%)	(1.5)	(0.7)	(6.6)	(8.0)	(21.9)	(13.1)	(16.8)	(1.5)	(10.2)	(5.8)	(4.4)	(4.4)	(3.6)		(1.5)	(100.0)				
210～	2	3	6	15	12	18	7	2	12	8	8	1	1		3	98		98	5.43	1.41
(%)	(2.0)	(3.1)	(6.1)	(15.3)	(12.2)	(18.4)	(7.1)	(2.0)	(12.2)	(8.2)	(8.2)	(1.0)	(1.0)		(3.1)	(100.0)				
240～	1	1	3	14	16	18	11	3	11	2	2	1	1	2	2	88		88	5.39	1.28
(%)	(1.1)	(1.1)	(3.4)	(15.9)	(18.2)	(20.5)	(12.5)	(3.4)	(12.5)	(2.3)	(2.3)	(1.1)	(1.1)	(2.3)	(2.3)	(100.0)				
300～		2	1	5	3	4	8		5	3	5	2		2	2	42		42	6.07	1.79
(%)		(4.8)	(2.4)	(11.9)	(7.1)	(9.5)	(19.0)		(11.9)	(7.1)	(11.9)	(4.8)		(4.8)	(4.8)	(100.0)				
350～	1	1	1	1	3	5	4		4	4	2	1		1		29		29	5.95	1.31
(%)	(3.4)		(3.4)	(3.4)	(10.3)	(17.2)	(13.8)		(13.8)	(13.8)	(6.9)	(3.4)		(3.4)		(100.0)				
400～	1	1	2	3	4	4	4		5	1	1				2	24		24	5.72	1.56
(%)	(4.2)		(4.2)	(8.3)	(12.5)	(16.7)	(16.7)		(20.8)	(4.2)	(4.2)				(8.3)	(100.0)				
500～	1	1	4	5	5	4	5		9	2	4	2	2	1		45		45	5.64	1.46
(%)	(2.2)	(2.2)	(8.9)	(11.1)	(11.1)	(8.9)	(11.1)		(20.0)	(4.4)	(8.9)	(4.4)	(4.4)	(2.2)		(100.0)				
合計	61	124	231	381	495	472	366	73	238	184	114	70	40	22	31	2,902	4	2,906	5.24	1.31
(%)	(2.1)	(4.3)	(8.0)	(13.1)	(17.1)	(16.3)	(12.6)	(2.5)	(8.2)	(6.3)	(3.9)	(2.4)	(1.4)	(0.8)	(1.1)	(100.0)				
記載なし				2	2	1	2		1		1	1	1			11	20	31	5.86	1.38
(%)				(18.2)	(18.2)	(9.1)	(18.2)		(9.1)		(9.1)	(9.1)	(9.1)			(100.0)				
総計	61	124	231	383	497	473	368	73	239	184	115	71	41	22	31	2,913	24	2,937	5.25	1.31
(%)	(2.1)	(4.3)	(7.9)	(13.1)	(17.1)	(16.2)	(12.6)	(2.5)	(8.2)	(6.3)	(3.9)	(2.4)	(1.4)	(0.8)	(1.1)	(100.0)				
平均	97.75	78.41	91.29	109.41	101.86	106.93	120.07	95.74	144.21	115.05	136.63	131.27	151.82	136.36	156.03	111.80	103.25	111.79		
標準偏差	103.71	74.71	90.36	114.72	92.97	98.10	121.16	64.78	127.15	94.25	138.72	123.48	154.01	137.69	120.12	108.19	30.84	108.12		

数値下のカッコ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査