

表 3478 透析前リン濃度(mg/dL) 透析前ヘモグロビン濃度(g/dL)別(血液透析濾過患者 透析回数週3回)

透析前ヘモグロビン濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
5未満		4		3	7	6	4				1				1	26		26	4.93	1.35
(%)		(15.4)		(11.5)	(26.9)	(23.1)	(15.4)				(3.8)				(3.8)	(100.0)				
5～		1	1	6	6	6	4	1		2	1	1		1	1	31		31	5.37	1.45
(%)		(3.2)	(3.2)	(19.4)	(19.4)	(19.4)	(12.9)	(3.2)		(6.5)	(3.2)	(3.2)		(3.2)	(3.2)	(100.0)				
6～	11	9	12	15	15	15	13	4	6	7	3	1	1	1	1	113	1	114	4.73	1.47
(%)	(9.7)	(8.0)	(10.6)	(13.3)	(13.3)	(13.3)	(11.5)	(3.5)	(5.3)	(6.2)	(2.7)	(0.9)	(0.9)	(0.9)	(100.0)					
7～	47	30	35	48	55	60	52	8	32	25	16	15	6	5	9	443	2	445	5.02	1.66
(%)	(10.6)	(6.8)	(7.9)	(10.8)	(12.4)	(13.5)	(11.7)	(1.8)	(7.2)	(5.6)	(3.6)	(3.4)	(1.4)	(1.1)	(2.0)	(100.0)				
8～	120	111	169	200	211	223	194	39	102	82	76	36	28	14	21	1,626	11	1,637	5.02	1.54
(%)	(7.4)	(6.8)	(10.4)	(12.3)	(13.0)	(13.7)	(11.9)	(2.4)	(6.3)	(5.0)	(4.7)	(2.2)	(1.7)	(0.9)	(1.3)	(100.0)				
9～	229	300	533	758	919	908	704	128	405	333	236	140	91	53	100	5,837	13	5,850	5.18	1.45
(%)	(3.9)	(5.1)	(9.1)	(13.0)	(15.7)	(15.6)	(12.1)	(2.2)	(6.9)	(5.7)	(4.0)	(2.4)	(1.6)	(0.9)	(1.7)	(100.0)				
10～	367	538	967	1,580	1,992	2,095	1,746	335	1,050	871	517	341	199	130	187	12,915	32	12,947	5.30	1.38
(%)	(2.8)	(4.2)	(7.5)	(12.2)	(15.4)	(16.2)	(13.5)	(2.6)	(8.1)	(6.7)	(4.0)	(2.6)	(1.5)	(1.0)	(1.4)	(100.0)				
11～	212	399	774	1,327	1,738	1,838	1,662	270	1,037	918	538	334	177	123	190	11,537	40	11,577	5.42	1.37
(%)	(1.8)	(3.5)	(6.7)	(11.5)	(15.1)	(15.9)	(14.4)	(2.3)	(9.0)	(8.0)	(4.7)	(2.9)	(1.5)	(1.1)	(1.6)	(100.0)				
12～	97	116	238	445	544	688	642	114	416	383	262	159	98	58	104	4,364	18	4,382	5.60	1.44
(%)	(2.2)	(2.7)	(5.5)	(10.2)	(12.5)	(15.8)	(14.7)	(2.6)	(9.5)	(8.8)	(6.0)	(3.6)	(2.2)	(1.3)	(2.4)	(100.0)				
13～	27	35	71	91	135	169	168	25	119	124	76	45	29	24	44	1,182	3	1,185	5.75	1.55
(%)	(2.3)	(3.0)	(6.0)	(7.7)	(11.4)	(14.3)	(14.2)	(2.1)	(10.1)	(10.5)	(6.4)	(3.8)	(2.5)	(2.0)	(3.7)	(100.0)				
14～	11	13	24	58	47	65	74	18	51	45	39	28	18	13	18	522	1	523	5.84	1.60
(%)	(2.1)	(2.5)	(4.6)	(11.1)	(9.0)	(12.5)	(14.2)	(3.4)	(9.8)	(8.6)	(7.5)	(5.4)	(3.4)	(2.5)	(3.4)	(100.0)				
合計	1,121	1,556	2,824	4,531	5,669	6,073	5,263	942	3,218	2,790	1,765	1,100	647	422	675	38,596	121	38,717	5.36	1.42
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.7)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.9)	(1.7)	(1.1)	(1.7)	(100.0)				
記載なし	12	18	25	34	43	61	36	8	26	16	10	4	4	1	7	305	924	1,229	5.19	1.52
(%)	(3.9)	(5.9)	(8.2)	(11.1)	(14.1)	(20.0)	(11.8)	(2.6)	(8.5)	(5.2)	(3.3)	(1.3)	(1.3)	(0.3)	(2.3)	(100.0)				
総計	1,133	1,574	2,849	4,565	5,712	6,134	5,299	950	3,244	2,806	1,775	1,104	651	423	682	38,901	1,045	39,946	5.36	1.43
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.8)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.8)	(1.7)	(1.1)	(1.8)	(100.0)				
平均	10.35	10.51	10.65	10.75	10.77	10.84	10.91	10.88	10.99	11.03	11.02	11.04	11.02	11.08	11.10	10.84	10.82	10.84		
標準偏差	1.44	1.28	1.23	1.25	1.19	1.22	1.23	1.30	1.22	1.24	1.33	1.37	1.34	1.42	1.44	1.26	1.32	1.26		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査