

表 3480 透析前リン濃度(mg/dL) HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

HDL-C濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
20未満	14	3	5	18	11	11	6		2	1	2	2	1	1	4	81		81	4.62	2.01
(%)	(17.3)	(3.7)	(6.2)	(22.2)	(13.6)	(13.6)	(7.4)		(2.5)	(1.2)	(2.5)	(2.5)	(1.2)	(1.2)	(4.9)	(100.0)				
20～	68	107	162	227	268	311	228	45	129	120	72	52	33	14	36	1,872	2	1,874	5.23	1.47
(%)	(3.6)	(5.7)	(8.7)	(12.1)	(14.3)	(16.6)	(12.2)	(2.4)	(6.9)	(6.4)	(3.8)	(2.8)	(1.8)	(0.7)	(1.9)	(100.0)				
30～	192	280	510	738	882	943	871	129	469	413	273	154	100	65	105	6,124	7	6,131	5.29	1.43
(%)	(3.1)	(4.6)	(8.3)	(12.1)	(14.4)	(15.4)	(14.2)	(2.1)	(7.7)	(6.7)	(4.5)	(2.5)	(1.6)	(1.1)	(1.7)	(100.0)				
40～	205	321	553	907	1,113	1,102	1,021	159	593	522	334	218	115	78	112	7,353	7	7,360	5.33	1.41
(%)	(2.8)	(4.4)	(7.5)	(12.3)	(15.1)	(15.0)	(13.9)	(2.2)	(8.1)	(7.1)	(4.5)	(3.0)	(1.6)	(1.1)	(1.5)	(100.0)				
50～	148	236	434	702	886	994	802	141	548	430	247	163	101	58	78	5,968	3	5,971	5.34	1.36
(%)	(2.5)	(4.0)	(7.3)	(11.8)	(14.8)	(16.7)	(13.4)	(2.4)	(9.2)	(7.2)	(4.1)	(2.7)	(1.7)	(1.0)	(1.3)	(100.0)				
60～	78	121	276	452	595	654	532	106	289	314	169	92	63	35	79	3,855	2	3,857	5.40	1.38
(%)	(2.0)	(3.1)	(7.2)	(11.7)	(15.4)	(17.0)	(13.8)	(2.7)	(7.5)	(8.1)	(4.4)	(2.4)	(1.6)	(0.9)	(2.0)	(100.0)				
70～	35	72	140	217	295	322	284	58	173	155	103	64	45	19	40	2,022	1	2,023	5.48	1.43
(%)	(1.7)	(3.6)	(6.9)	(10.7)	(14.6)	(15.9)	(14.0)	(2.9)	(8.6)	(7.7)	(5.1)	(3.2)	(2.2)	(0.9)	(2.0)	(100.0)				
80～	26	43	47	108	154	163	157	26	102	70	50	29	8	10	20	1,013		1,013	5.42	1.37
(%)	(2.6)	(4.2)	(4.6)	(10.7)	(15.2)	(16.1)	(15.5)	(2.6)	(10.1)	(6.9)	(4.9)	(2.9)	(0.8)	(1.0)	(2.0)	(100.0)				
90～	16	10	26	53	62	61	63	13	38	36	16	17	7	5	8	431		431	5.44	1.43
(%)	(3.7)	(2.3)	(6.0)	(12.3)	(14.4)	(14.2)	(14.6)	(3.0)	(8.8)	(8.4)	(3.7)	(3.9)	(1.6)	(1.2)	(1.9)	(100.0)				
100～	7	9	21	38	57	58	55	12	38	35	19	14	10	2	7	382	1	383	5.56	1.36
(%)	(1.8)	(2.4)	(5.5)	(9.9)	(14.9)	(15.2)	(14.4)	(3.1)	(9.9)	(9.2)	(5.0)	(3.7)	(2.6)	(0.5)	(1.8)	(100.0)				
合計	789	1,202	2,174	3,460	4,323	4,619	4,019	689	2,381	2,096	1,285	805	483	287	489	29,101	23	29,124	5.34	1.41
(%)	(2.7)	(4.1)	(7.5)	(11.9)	(14.9)	(15.9)	(13.8)	(2.4)	(8.2)	(7.2)	(4.4)	(2.8)	(1.7)	(1.0)	(1.7)	(100.0)				
記載なし	344	372	675	1,105	1,389	1,515	1,280	261	863	710	490	299	168	136	193	9,800	1,022	10,822	5.39	1.48
(%)	(3.5)	(3.8)	(6.9)	(11.3)	(14.2)	(15.5)	(13.1)	(2.7)	(8.8)	(7.2)	(5.0)	(3.1)	(1.7)	(1.4)	(2.0)	(100.0)				
総計	1,133	1,574	2,849	4,565	5,712	6,134	5,299	950	3,244	2,806	1,775	1,104	651	423	682	38,901	1,045	39,946	5.36	1.43
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.8)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.8)	(1.7)	(1.1)	(1.8)	(100.0)				
平均	48.06	48.43	49.13	50.21	51.04	50.99	51.24	52.72	51.94	51.99	51.42	51.79	51.61	50.57	51.37	50.86	46.13	50.85		
標準偏差	17.56	16.42	16.13	16.87	17.22	16.96	17.05	18.27	17.44	17.63	17.19	18.41	18.71	16.36	18.34	17.18	17.53	17.18		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査