

表 3481 透析前リン濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

Non-HDL-C濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
50未満	31	36	45	74	78	69	53	11	34	41	26	13	8	4	6	529		529	5.09	1.47
(%)	(5.9)	(6.8)	(8.5)	(14.0)	(14.7)	(13.0)	(10.0)	(2.1)	(6.4)	(7.8)	(4.9)	(2.5)	(1.5)	(0.8)	(1.1)	(100.0)				
50～	101	109	201	312	358	346	325	51	182	171	101	56	36	18	35	2,402	1	2,403	5.21	1.43
(%)	(4.2)	(4.5)	(8.4)	(13.0)	(14.9)	(14.4)	(13.5)	(2.1)	(7.6)	(7.1)	(4.2)	(2.3)	(1.5)	(0.7)	(1.5)	(100.0)				
70～	176	223	380	636	783	811	727	101	402	343	209	150	81	36	73	5,131	7	5,138	5.27	1.40
(%)	(3.4)	(4.3)	(7.4)	(12.4)	(15.3)	(15.8)	(14.2)	(2.0)	(7.8)	(6.7)	(4.1)	(2.9)	(1.6)	(0.7)	(1.4)	(100.0)				
90～	159	273	498	779	952	979	819	146	529	431	274	148	107	73	94	6,261	4	6,265	5.31	1.40
(%)	(2.5)	(4.4)	(8.0)	(12.4)	(15.2)	(15.6)	(13.1)	(2.3)	(8.4)	(6.9)	(4.4)	(2.4)	(1.7)	(1.2)	(1.5)	(100.0)				
110～	113	188	379	600	775	833	699	128	446	379	229	152	85	52	102	5,160	2	5,162	5.40	1.40
(%)	(2.2)	(3.6)	(7.3)	(11.6)	(15.0)	(16.1)	(13.5)	(2.5)	(8.6)	(7.3)	(4.4)	(2.9)	(1.6)	(1.0)	(2.0)	(100.0)				
130～	62	118	221	341	456	545	451	75	260	264	132	76	66	24	54	3,145	4	3,149	5.40	1.37
(%)	(2.0)	(3.8)	(7.0)	(10.8)	(14.5)	(17.3)	(14.3)	(2.4)	(8.3)	(8.4)	(4.2)	(2.4)	(2.1)	(0.8)	(1.7)	(100.0)				
150～	28	58	91	167	217	243	249	44	125	134	94	63	24	25	35	1,597		1,597	5.55	1.43
(%)	(1.8)	(3.6)	(5.7)	(10.5)	(13.6)	(15.2)	(15.6)	(2.8)	(7.8)	(8.4)	(5.9)	(3.9)	(1.5)	(1.6)	(2.2)	(100.0)				
170～	19	17	52	75	78	94	111	23	52	64	36	17	16	13	15	682		682	5.53	1.47
(%)	(2.8)	(2.5)	(7.6)	(11.0)	(11.4)	(13.8)	(16.3)	(3.4)	(7.6)	(9.4)	(5.3)	(2.5)	(2.3)	(1.9)	(2.2)	(100.0)				
190～	1	7	18	19	32	35	44	12	22	20	14	14	8	7	9	262		262	5.81	1.49
(%)	(0.4)	(2.7)	(6.9)	(7.3)	(12.2)	(13.4)	(16.8)	(4.6)	(8.4)	(7.6)	(5.3)	(5.3)	(3.1)	(2.7)	(3.4)	(100.0)				
210～	3	5	12	13	19	20	14	2	13	9	8	6	3	5	9	141		141	5.77	1.80
(%)	(2.1)	(3.5)	(8.5)	(9.2)	(13.5)	(14.2)	(9.9)	(1.4)	(9.2)	(6.4)	(5.7)	(4.3)	(2.1)	(3.5)	(6.4)	(100.0)				
合計	693	1,034	1,897	3,016	3,748	3,975	3,492	593	2,065	1,856	1,123	695	434	257	432	25,310	18	25,328	5.35	1.41
(%)	(2.7)	(4.1)	(7.5)	(11.9)	(14.8)	(15.7)	(13.8)	(2.3)	(8.2)	(7.3)	(4.4)	(2.7)	(1.7)	(1.0)	(1.7)	(100.0)				
記載なし	440	540	952	1,549	1,964	2,159	1,807	357	1,179	950	652	409	217	166	250	13,591	1,027	14,618	5.37	1.45
(%)	(3.2)	(4.0)	(7.0)	(11.4)	(14.5)	(15.9)	(13.3)	(2.6)	(8.7)	(7.0)	(4.8)	(3.0)	(1.6)	(1.2)	(1.8)	(100.0)				
総計	1,133	1,574	2,849	4,565	5,712	6,134	5,299	950	3,244	2,806	1,775	1,104	651	423	682	38,901	1,045	39,946	5.36	1.43
(%)	(2.9)	(4.0)	(7.3)	(11.7)	(14.7)	(15.8)	(13.6)	(2.4)	(8.3)	(7.2)	(4.6)	(2.8)	(1.7)	(1.1)	(1.8)	(100.0)				
平均	97.92	103.04	105.35	104.73	105.74	107.68	108.62	111.15	107.95	109.26	109.22	109.79	110.30	115.93	114.07	107.12	100.83	107.12		
標準偏差	32.95	32.68	32.99	32.71	31.99	32.79	33.48	34.43	32.67	34.64	34.91	35.39	34.31	38.04	37.07	33.38	22.46	33.37		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査