

表 3496 リン濃度(mg/dL) Non-HDL-C濃度(mg/dL)別（腹膜透析患者）

Non-HDL-C濃度	3未満	3.0～	3.5～	4.0～	4.5～	5.0～	5.5～	6.0～	6.1～	6.5～	7.0～	7.5～	8.0～	8.5～	9.0～	合計	記載なし	総計	平均	標準偏差
50未満	1	1	1	3	2	3	2		1	1	1	2				18		18	5.24	1.48
(%)	(5.6)	(5.6)	(5.6)	(16.7)	(11.1)	(16.7)	(11.1)		(5.6)	(5.6)	(5.6)	(11.1)				(100.0)				
50～	1	5	11	12	18	20	15		15	7	3	1	1	2	2	113	1	114	5.31	1.31
(%)	(0.9)	(4.4)	(9.7)	(10.6)	(15.9)	(17.7)	(13.3)		(13.3)	(6.2)	(2.7)	(0.9)	(0.9)	(1.8)	(1.8)	(100.0)				
70～	11	11	28	47	56	52	50	6	29	21	13	11	5	5	2	347	4	351	5.24	1.34
(%)	(3.2)	(3.2)	(8.1)	(13.5)	(16.1)	(15.0)	(14.4)	(1.7)	(8.4)	(6.1)	(3.7)	(3.2)	(1.4)	(1.4)	(0.6)	(100.0)				
90～	21	34	50	79	102	97	91	16	57	43	27	19	8	3	7	654	6	660	5.22	1.35
(%)	(3.2)	(5.2)	(7.6)	(12.1)	(15.6)	(14.8)	(13.9)	(2.4)	(8.7)	(6.6)	(4.1)	(2.9)	(1.2)	(0.5)	(1.1)	(100.0)				
110～	12	22	81	117	119	122	108	21	53	60	30	24	10	3	6	788	9	797	5.24	1.29
(%)	(1.5)	(2.8)	(10.3)	(14.8)	(15.1)	(15.5)	(13.7)	(2.7)	(6.7)	(7.6)	(3.8)	(3.0)	(1.3)	(0.4)	(0.8)	(100.0)				
130～	9	23	54	84	103	115	81	20	63	45	20	15	9	3	9	653	9	662	5.30	1.30
(%)	(1.4)	(3.5)	(8.3)	(12.9)	(15.8)	(17.6)	(12.4)	(3.1)	(9.6)	(6.9)	(3.1)	(2.3)	(1.4)	(0.5)	(1.4)	(100.0)				
150～	5	18	37	61	65	72	65	11	33	18	19	13	4	2	10	433	3	436	5.29	1.39
(%)	(1.2)	(4.2)	(8.5)	(14.1)	(15.0)	(16.6)	(15.0)	(2.5)	(7.6)	(4.2)	(4.4)	(3.0)	(0.9)	(0.5)	(2.3)	(100.0)				
170～	6	12	26	31	35	41	50	10	9	18	12	4	1	2	5	262	1	263	5.23	1.36
(%)	(2.3)	(4.6)	(9.9)	(11.8)	(13.4)	(15.6)	(19.1)	(3.8)	(3.4)	(6.9)	(4.6)	(1.5)	(0.4)	(0.8)	(1.9)	(100.0)				
190～	1	7	9	12	15	25	13	3	14	13	3	1	1	1	2	120		120	5.34	1.28
(%)	(0.8)	(5.8)	(7.5)	(10.0)	(12.5)	(20.8)	(10.8)	(2.5)	(11.7)	(10.8)	(2.5)	(0.8)	(0.8)	(0.8)	(1.7)	(100.0)				
210～	2	4	11	4	19	9	15	2	10	5	1	3	1	1	1	88	2	90	5.22	1.42
(%)	(2.3)	(4.5)	(12.5)	(4.5)	(21.6)	(10.2)	(17.0)	(2.3)	(11.4)	(5.7)	(1.1)	(3.4)	(1.1)	(1.1)	(1.1)	(100.0)				
合計	69	137	308	450	534	556	490	89	284	231	129	93	40	22	44	3,476	35	3,511	5.26	1.33
(%)	(2.0)	(3.9)	(8.9)	(12.9)	(15.4)	(16.0)	(14.1)	(2.6)	(8.2)	(6.6)	(3.7)	(2.7)	(1.2)	(0.6)	(1.3)	(100.0)				
記載なし	57	133	217	320	387	392	349	49	193	168	83	56	31	16	45	2,496	2,934	5,430	5.23	1.38
(%)	(2.3)	(5.3)	(8.7)	(12.8)	(15.5)	(15.7)	(14.0)	(2.0)	(7.7)	(6.7)	(3.3)	(2.2)	(1.2)	(0.6)	(1.8)	(100.0)				
総計	126	270	525	770	921	948	839	138	477	399	212	149	71	38	89	5,972	2,969	8,941	5.25	1.35
(%)	(2.1)	(4.5)	(8.8)	(12.9)	(15.4)	(15.9)	(14.0)	(2.3)	(8.0)	(6.7)	(3.5)	(2.5)	(1.2)	(0.6)	(1.5)	(100.0)				
平均	118.93	130.64	130.02	126.37	128.91	129.08	129.81	135.28	127.65	128.37	125.54	124.06	124.15	121.86	137.36	128.48	127.37	128.47		
標準偏差	37.55	40.58	38.31	33.77	40.21	37.57	37.95	36.09	40.78	38.31	34.66	37.33	34.46	45.36	38.46	37.98	41.60	38.01		

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2014年調査