

表 3501 PTH値(pg/mL) 治療方法(新分類)別(透析患者全体) PTH測定法:intact-PTH

治療方法 (新分類)	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
施設血液透析	9,872	14,270	15,116	16,113	16,261	15,854	15,147	14,035	12,598	28,506	351	16,254	9,366	5,369	3,155	770	1,258	1,360	2,228	1,684	199,567	5,018	204,585	170.17	161.05
(%)	(4.9)	(7.2)	(7.6)	(8.1)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.1)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
Off-line HDF	322	348	448	408	420	441	404	335	300	689	14	407	248	148	80	26	32	39	66	63	5,238	109	5,347	172.21	177.86
(%)	(6.1)	(6.6)	(8.6)	(7.8)	(8.0)	(8.4)	(7.7)	(6.4)	(5.7)	(13.2)	(0.3)	(7.8)	(4.7)	(2.8)	(1.5)	(0.5)	(0.6)	(0.7)	(1.3)	(1.2)	(100.0)				
On-line HDF	1,372	1,861	2,000	2,195	2,442	2,367	2,184	1,998	1,864	4,337	64	2,565	1,475	905	514	118	205	218	369	285	29,338	768	30,106	178.29	167.88
(%)	(4.7)	(6.3)	(6.8)	(7.5)	(8.3)	(8.1)	(7.4)	(6.8)	(6.4)	(14.8)	(0.2)	(8.7)	(5.0)	(3.1)	(1.8)	(0.4)	(0.7)	(0.7)	(1.3)	(1.0)	(100.0)				
Push/Pull HDF	19	32	25	36	32	34	36	34	26	73	2	32	18	15	6	2	2	1	2	4	431	47	478	168.07	149.34
(%)	(4.4)	(7.4)	(5.8)	(8.4)	(7.4)	(7.9)	(8.4)	(7.9)	(6.0)	(16.9)	(0.5)	(7.4)	(4.2)	(3.5)	(1.4)	(0.5)	(0.5)	(0.2)	(0.5)	(0.9)	(100.0)				
ハイオフィルトレーション	12	12	19	25	17	16	24	14	15	35	1	24	16	7	8	2	4	2	4	4	261	3	264	192.53	169.78
(%)	(4.6)	(4.6)	(7.3)	(9.6)	(6.5)	(6.1)	(9.2)	(5.4)	(5.7)	(13.4)	(0.4)	(9.2)	(6.1)	(2.7)	(3.1)	(0.8)	(1.5)	(0.8)	(1.5)	(1.5)	(100.0)				
血液濾過	17	17	8	8	1	8	3	2		4		3	1	4	1			2	3	2	84	5	89	149.43	218.15
(%)	(20.2)	(20.2)	(9.5)	(9.5)	(1.2)	(9.5)	(3.6)	(2.4)		(4.8)		(3.6)	(1.2)	(4.8)	(1.2)			(2.4)	(3.6)	(2.4)	(100.0)				
血液吸着透析 (リクセル)	127	97	115	108	109	103	99	81	72	168	2	96	53	30	14	3	4	11	16	2	1,310	21	1,331	148.43	132.42
(%)	(9.7)	(7.4)	(8.8)	(8.2)	(8.3)	(7.9)	(7.6)	(6.2)	(5.5)	(12.8)	(0.2)	(7.3)	(4.0)	(2.3)	(1.1)	(0.2)	(0.3)	(0.8)	(1.2)	(0.2)	(100.0)				
血液吸着透析 (その他)					1							1						1			3		3	315.67	255.15
(%)					(33.3)							(33.3)						(33.3)			(100.0)				
在宅血液透析	9	28	17	22	21	19	20	23	22	47		26	18	6	8		5	7	6	8	312	37	349	203.87	184.85
(%)	(2.9)	(9.0)	(5.4)	(7.1)	(6.7)	(6.1)	(6.4)	(7.4)	(7.1)	(15.1)		(8.3)	(5.8)	(1.9)	(2.6)		(1.6)	(2.2)	(1.9)	(2.6)	(100.0)				
腹膜透析 (CAPD)	62	168	151	174	173	185	164	157	149	338	7	242	168	100	85	22	28	43	54	52	2,522	203	2,725	216.43	202.69
(%)	(2.5)	(6.7)	(6.0)	(6.9)	(6.9)	(7.3)	(6.5)	(6.2)	(5.9)	(13.4)	(0.3)	(9.6)	(6.7)	(4.0)	(3.4)	(0.9)	(1.1)	(1.7)	(2.1)	(2.1)	(100.0)				
腹膜透析 (APD)	44	78	79	100	90	94	91	91	86	211	2	139	121	65	36	17	21	21	34	48	1,468	143	1,611	240.80	281.13
(%)	(3.0)	(5.3)	(5.4)	(6.8)	(6.1)	(6.4)	(6.2)	(6.2)	(5.9)	(14.4)	(0.1)	(9.5)	(8.2)	(4.4)	(2.5)	(1.2)	(1.4)	(1.4)	(2.3)	(3.3)	(100.0)				
腹膜透析 (CCPD)	17	35	38	41	46	37	50	37	41	105	1	50	42	35	18	4	5	9	17	10	638	30	668	214.36	190.69
(%)	(2.7)	(5.5)	(6.0)	(6.4)	(7.2)	(5.8)	(7.8)	(5.8)	(6.4)	(16.5)	(0.2)	(7.8)	(6.6)	(5.5)	(2.8)	(0.6)	(0.8)	(1.4)	(2.7)	(1.6)	(100.0)				
合計	11,873	16,946	18,016	19,230	19,613	19,158	18,222	16,807	15,173	34,513	444	19,839	11,526	6,684	3,925	964	1,564	1,714	2,799	2,162	241,172	6,384	247,556	172.17	163.94
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
記載なし																									
(%)																									
総計	11,873	16,946	18,016	19,230	19,613	19,158	18,222	16,807	15,173	34,513	444	19,839	11,526	6,684	3,925	964	1,564	1,714	2,799	2,162	241,172	6,384	247,556	172.17	163.94
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの
(C)Japanese Society for Dialysis Therapy
2014年調査