

表 3510 PTH値(pg/mL) 総コレステロール濃度(mg/dL)別 (透析患者全体) PTH測定法:intact-PTH

総コレステロール濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
100未満	478	577	634	638	609	582	524	473	414	870	15	457	230	130	84	17	31	20	42	34	6,859	155	7,014	144.82	132.51
(%)	(7.0)	(8.4)	(9.2)	(9.3)	(8.9)	(8.5)	(7.6)	(6.9)	(6.0)	(12.7)	(0.2)	(6.7)	(3.4)	(1.9)	(1.2)	(0.2)	(0.5)	(0.3)	(0.6)	(0.5)	(100.0)				
100～	1,060	1,550	1,680	1,684	1,764	1,681	1,514	1,393	1,224	2,734	32	1,532	847	446	241	53	105	117	151	105	19,913	314	20,227	156.83	142.80
(%)	(5.3)	(7.8)	(8.4)	(8.5)	(8.9)	(8.4)	(7.6)	(7.0)	(6.1)	(13.7)	(0.2)	(7.7)	(4.3)	(2.2)	(1.2)	(0.3)	(0.5)	(0.6)	(0.8)	(0.5)	(100.0)				
120～	1,882	2,706	2,912	3,044	3,160	3,067	2,863	2,713	2,360	5,240	52	3,062	1,634	914	519	128	210	239	340	259	37,304	554	37,858	164.28	150.68
(%)	(5.0)	(7.3)	(7.8)	(8.2)	(8.5)	(8.2)	(7.7)	(7.3)	(6.3)	(14.0)	(0.1)	(8.2)	(4.4)	(2.5)	(1.4)	(0.3)	(0.6)	(0.6)	(0.9)	(0.7)	(100.0)				
140～	2,027	3,002	3,049	3,411	3,450	3,480	3,354	2,997	2,844	6,232	63	3,553	2,074	1,165	649	181	285	284	459	323	42,882	607	43,489	170.59	156.74
(%)	(4.7)	(7.0)	(7.1)	(8.0)	(8.0)	(8.1)	(7.8)	(7.0)	(6.6)	(14.5)	(0.1)	(8.3)	(4.8)	(2.7)	(1.5)	(0.4)	(0.7)	(0.7)	(1.1)	(0.8)	(100.0)				
160～	1,657	2,343	2,575	2,693	2,824	2,750	2,635	2,441	2,304	5,161	86	2,997	1,794	1,044	622	141	247	249	442	339	35,344	513	35,857	176.89	167.25
(%)	(4.7)	(6.6)	(7.3)	(7.6)	(8.0)	(7.8)	(7.5)	(6.9)	(6.5)	(14.6)	(0.2)	(8.5)	(5.1)	(3.0)	(1.8)	(0.4)	(0.7)	(0.7)	(1.3)	(1.0)	(100.0)				
180～	1,024	1,526	1,664	1,685	1,746	1,754	1,659	1,543	1,392	3,291	44	1,905	1,121	694	447	102	156	178	312	250	22,493	299	22,792	180.36	172.05
(%)	(4.6)	(6.8)	(7.4)	(7.5)	(7.8)	(7.8)	(7.4)	(6.9)	(6.2)	(14.6)	(0.2)	(8.5)	(5.0)	(3.1)	(2.0)	(0.5)	(0.7)	(0.8)	(1.4)	(1.1)	(100.0)				
200～	555	742	841	944	907	926	868	792	723	1,754	18	1,072	689	381	252	63	94	99	183	131	12,034	165	12,199	185.17	179.03
(%)	(4.6)	(6.2)	(7.0)	(7.8)	(7.5)	(7.7)	(7.2)	(6.6)	(6.0)	(14.6)	(0.1)	(8.9)	(5.7)	(3.2)	(2.1)	(0.5)	(0.8)	(0.8)	(1.5)	(1.1)	(100.0)				
220～	255	356	343	409	396	424	352	360	317	719	16	471	323	175	92	31	41	41	86	74	5,281	82	5,363	188.82	194.65
(%)	(4.8)	(6.7)	(6.5)	(7.7)	(7.5)	(8.0)	(6.7)	(6.8)	(6.0)	(13.6)	(0.3)	(8.9)	(6.1)	(3.3)	(1.7)	(0.6)	(0.8)	(0.8)	(1.6)	(1.4)	(100.0)				
240～	111	154	142	160	164	128	137	137	115	296	7	203	121	72	38	8	10	29	37	24	2,093	25	2,118	188.05	190.33
(%)	(5.3)	(7.4)	(6.8)	(7.6)	(7.8)	(6.1)	(6.5)	(6.5)	(5.5)	(14.1)	(0.3)	(9.7)	(5.8)	(3.4)	(1.8)	(0.4)	(0.5)	(1.4)	(1.8)	(1.1)	(100.0)				
260～	68	93	93	111	86	102	95	83	77	175	3	115	76	61	40	8	16	23	23	30	1,378	24	1,402	208.05	244.05
(%)	(4.9)	(6.7)	(6.7)	(8.1)	(6.2)	(7.4)	(6.9)	(6.0)	(5.6)	(12.7)	(0.2)	(8.3)	(5.5)	(4.4)	(2.9)	(0.6)	(1.2)	(1.7)	(1.7)	(2.2)	(100.0)				
合計	9,117	13,049	13,933	14,779	15,106	14,894	14,001	12,932	11,770	26,472	336	15,367	8,909	5,082	2,984	732	1,195	1,279	2,075	1,569	185,581	2,738	188,319	171.22	161.58
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(8.0)	(7.5)	(7.0)	(6.3)	(14.3)	(0.2)	(8.3)	(4.8)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
記載なし	2,756	3,897	4,083	4,451	4,507	4,264	4,221	3,875	3,403	8,041	108	4,472	2,617	1,602	941	232	369	435	724	593	55,591	3,646	59,237	175.37	171.57
(%)	(5.0)	(7.0)	(7.3)	(8.0)	(8.1)	(7.7)	(7.6)	(7.0)	(6.1)	(14.5)	(0.2)	(8.0)	(4.7)	(2.9)	(1.7)	(0.4)	(0.7)	(0.8)	(1.3)	(1.1)	(100.0)				
総計	11,873	16,946	18,016	19,230	19,613	19,158	18,222	16,807	15,173	34,513	444	19,839	11,526	6,684	3,925	964	1,564	1,714	2,799	2,162	241,172	6,384	247,556	172.17	163.94
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
平均	153.47	153.80	153.62	154.42	153.91	154.60	155.00	155.12	155.46	156.30	160.34	157.46	159.65	160.73	161.81	162.30	160.34	162.67	164.19	165.68	155.80	153.32	155.77		
標準偏差	36.73	36.15	36.02	35.97	35.18	35.12	35.03	35.24	34.77	34.91	37.31	35.60	35.84	36.48	36.46	36.32	35.78	37.58	36.23	37.96	35.65	37.08	35.67		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査