

表 3511 PTH値(pg/mL) HDL-C濃度(mg/dL)別 (透析患者全体) PTH測定法:intact-PTH

HDL-C濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
20未満	86	82	78	54	56	45	39	41	31	61	2	53	26	10	9	4	2		5	4	688	21	709	132.38	133.78
(%)	(12.5)	(11.9)	(11.3)	(7.8)	(8.1)	(6.5)	(5.7)	(6.0)	(4.5)	(8.9)	(0.3)	(7.7)	(3.8)	(1.5)	(1.3)	(0.6)	(0.3)		(0.7)	(0.6)	(100.0)				
20～	763	1,019	1,060	1,120	1,114	1,109	1,013	965	876	1,948	27	1,143	610	317	189	52	80	93	133	71	13,702	213	13,915	162.54	141.49
(%)	(5.6)	(7.4)	(7.7)	(8.2)	(8.1)	(8.1)	(7.4)	(7.0)	(6.4)	(14.2)	(0.2)	(8.3)	(4.5)	(2.3)	(1.4)	(0.4)	(0.6)	(0.7)	(1.0)	(0.5)	(100.0)				
30～	1,997	2,878	2,960	3,192	3,201	3,330	3,110	2,808	2,562	5,800	68	3,367	1,896	1,095	612	164	260	297	405	326	40,328	601	40,929	169.68	154.03
(%)	(5.0)	(7.1)	(7.3)	(7.9)	(7.9)	(8.3)	(7.7)	(7.0)	(6.4)	(14.4)	(0.2)	(8.3)	(4.7)	(2.7)	(1.5)	(0.4)	(0.6)	(0.7)	(1.0)	(0.8)	(100.0)				
40～	2,197	3,066	3,415	3,528	3,667	3,621	3,479	3,228	2,891	6,495	86	3,750	2,129	1,280	719	199	295	319	493	404	45,261	572	45,833	172.37	164.60
(%)	(4.9)	(6.8)	(7.5)	(7.8)	(8.1)	(8.0)	(7.7)	(7.1)	(6.4)	(14.4)	(0.2)	(8.3)	(4.7)	(2.8)	(1.6)	(0.4)	(0.7)	(0.7)	(1.1)	(0.9)	(100.0)				
50～	1,492	2,282	2,461	2,673	2,744	2,514	2,523	2,286	2,153	4,735	58	2,753	1,607	998	575	128	217	208	419	314	33,140	400	33,540	175.33	172.56
(%)	(4.5)	(6.9)	(7.4)	(8.1)	(8.3)	(7.6)	(7.6)	(6.9)	(6.5)	(14.3)	(0.2)	(8.3)	(4.8)	(3.0)	(1.7)	(0.4)	(0.7)	(0.6)	(1.3)	(0.9)	(100.0)				
60～	831	1,258	1,420	1,578	1,570	1,596	1,463	1,359	1,295	2,861	34	1,696	1,037	554	350	59	126	144	208	185	19,624	266	19,890	175.96	164.28
(%)	(4.2)	(6.4)	(7.2)	(8.0)	(8.0)	(8.1)	(7.5)	(6.9)	(6.6)	(14.6)	(0.2)	(8.6)	(5.3)	(2.8)	(1.8)	(0.3)	(0.6)	(0.7)	(1.1)	(0.9)	(100.0)				
70～	428	617	690	740	797	792	757	686	640	1,439	20	811	489	238	160	40	60	85	111	98	9,698	111	9,809	176.48	172.07
(%)	(4.4)	(6.4)	(7.1)	(7.6)	(8.2)	(8.2)	(7.8)	(7.1)	(6.6)	(14.8)	(0.2)	(8.4)	(5.0)	(2.5)	(1.6)	(0.4)	(0.6)	(0.9)	(1.1)	(1.0)	(100.0)				
80～	205	296	344	394	339	370	363	321	283	642	8	385	215	149	78	19	32	35	57	40	4,575	51	4,626	175.94	172.33
(%)	(4.5)	(6.5)	(7.5)	(8.6)	(7.4)	(8.1)	(7.9)	(7.0)	(6.2)	(14.0)	(0.2)	(8.4)	(4.7)	(3.3)	(1.7)	(0.4)	(0.7)	(0.8)	(1.2)	(0.9)	(100.0)				
90～	78	133	146	155	177	169	164	126	131	287	7	198	102	69	34	15	11	20	31	21	2,074	14	2,088	183.70	176.42
(%)	(3.8)	(6.4)	(7.0)	(7.5)	(8.5)	(8.1)	(7.9)	(6.1)	(6.3)	(13.8)	(0.3)	(9.5)	(4.9)	(3.3)	(1.6)	(0.7)	(0.5)	(1.0)	(1.5)	(1.0)	(100.0)				
100～	89	134	136	121	159	149	131	109	107	276	3	136	78	58	27	1	12	15	27	14	1,782	19	1,801	171.52	151.31
(%)	(5.0)	(7.5)	(7.6)	(6.8)	(8.9)	(8.4)	(7.4)	(6.1)	(6.0)	(15.5)	(0.2)	(7.6)	(4.4)	(3.3)	(1.5)	(0.1)	(0.7)	(0.8)	(1.5)	(0.8)	(100.0)				
合計	8,166	11,765	12,710	13,555	13,824	13,695	13,042	11,929	10,969	24,544	313	14,292	8,189	4,768	2,753	681	1,095	1,216	1,889	1,477	170,872	2,268	173,140	172.23	162.61
(%)	(4.8)	(6.9)	(7.4)	(7.9)	(8.1)	(8.0)	(7.6)	(7.0)	(6.4)	(14.4)	(0.2)	(8.4)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.9)	(100.0)				
記載なし	3,707	5,181	5,306	5,675	5,789	5,463	5,180	4,878	4,204	9,969	131	5,547	3,337	1,916	1,172	283	469	498	910	685	70,300	4,116	74,416	172.04	167.15
(%)	(5.3)	(7.4)	(7.5)	(8.1)	(8.2)	(7.8)	(7.4)	(6.9)	(6.0)	(14.2)	(0.2)	(7.9)	(4.7)	(2.7)	(1.7)	(0.4)	(0.7)	(0.7)	(1.3)	(1.0)	(100.0)				
総計	11,873	16,946	18,016	19,230	19,613	19,158	18,222	16,807	15,173	34,513	444	19,839	11,526	6,684	3,925	964	1,564	1,714	2,799	2,162	241,172	6,384	247,556	172.17	163.94
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
平均	47.53	48.17	48.57	48.67	48.88	48.76	48.89	48.62	48.92	48.90	49.28	48.91	49.28	49.54	49.64	47.93	49.08	49.46	50.22	50.38	48.79	46.91	48.76		
標準偏差	16.88	16.89	16.90	16.58	16.76	16.80	16.65	16.37	16.57	16.76	17.23	16.75	16.61	16.95	16.57	15.78	16.95	16.83	17.59	16.10	16.73	16.21	16.72		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査