

表 3512 PTH値(pg/mL) Non-HDL-C濃度(mg/dL)別 (透析患者全体) PTH測定法:intact-PTH

Non-HDL-C濃度	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
50未満	186	223	277	265	298	264	243	240	195	405	5	216	110	61	36	5	14	12	16	17	3,088	36	3,124	149.39	131.46
(%)	(6.0)	(7.2)	(9.0)	(8.6)	(9.7)	(8.5)	(7.9)	(7.8)	(6.3)	(13.1)	(0.2)	(7.0)	(3.6)	(2.0)	(1.2)	(0.2)	(0.5)	(0.4)	(0.5)	(0.6)	(100.0)				
50～	747	1,116	1,127	1,220	1,251	1,208	1,105	1,019	890	1,993	21	1,116	589	337	170	44	81	69	109	84	14,296	154	14,450	158.13	146.94
(%)	(5.2)	(7.8)	(7.9)	(8.5)	(8.8)	(8.4)	(7.7)	(7.1)	(6.2)	(13.9)	(0.1)	(7.8)	(4.1)	(2.4)	(1.2)	(0.3)	(0.6)	(0.5)	(0.8)	(0.6)	(100.0)				
70～	1,520	2,250	2,491	2,600	2,636	2,637	2,445	2,184	2,083	4,456	54	2,512	1,423	782	445	91	169	200	275	212	31,465	362	31,827	164.28	153.27
(%)	(4.8)	(7.2)	(7.9)	(8.3)	(8.4)	(8.4)	(7.8)	(6.9)	(6.6)	(14.2)	(0.2)	(8.0)	(4.5)	(2.5)	(1.4)	(0.3)	(0.5)	(0.6)	(0.9)	(0.7)	(100.0)				
90～	1,778	2,568	2,813	2,978	3,030	2,983	2,959	2,692	2,436	5,476	49	3,157	1,790	987	570	142	242	244	410	290	37,594	407	38,001	170.31	155.67
(%)	(4.7)	(6.8)	(7.5)	(7.9)	(8.1)	(7.9)	(7.9)	(7.2)	(6.5)	(14.6)	(0.1)	(8.4)	(4.8)	(2.6)	(1.5)	(0.4)	(0.6)	(0.6)	(1.1)	(0.8)	(100.0)				
110～	1,389	2,093	2,158	2,352	2,414	2,329	2,294	2,050	1,952	4,354	81	2,569	1,466	905	507	130	203	235	351	284	30,116	356	30,472	175.86	167.19
(%)	(4.6)	(6.9)	(7.2)	(7.8)	(8.0)	(7.7)	(7.6)	(6.8)	(6.5)	(14.5)	(0.3)	(8.5)	(4.9)	(3.0)	(1.7)	(0.4)	(0.7)	(0.8)	(1.2)	(0.9)	(100.0)				
130～	865	1,217	1,344	1,401	1,417	1,506	1,319	1,289	1,192	2,608	30	1,655	940	579	368	86	142	137	251	201	18,547	198	18,745	180.54	171.22
(%)	(4.7)	(6.6)	(7.2)	(7.6)	(7.6)	(8.1)	(7.1)	(6.9)	(6.4)	(14.1)	(0.2)	(8.9)	(5.1)	(3.1)	(2.0)	(0.5)	(0.8)	(0.7)	(1.4)	(1.1)	(100.0)				
150～	449	581	644	675	726	702	697	649	527	1,364	23	856	536	312	182	50	69	76	137	106	9,361	104	9,465	185.29	177.75
(%)	(4.8)	(6.2)	(6.9)	(7.2)	(7.8)	(7.5)	(7.4)	(6.9)	(5.6)	(14.6)	(0.2)	(9.1)	(5.7)	(3.3)	(1.9)	(0.5)	(0.7)	(0.8)	(1.5)	(1.1)	(100.0)				
170～	204	277	287	313	285	303	290	287	256	593	7	354	249	140	78	24	31	38	49	49	4,114	50	4,164	185.72	189.95
(%)	(5.0)	(6.7)	(7.0)	(7.6)	(6.9)	(7.4)	(7.0)	(7.0)	(6.2)	(14.4)	(0.2)	(8.6)	(6.1)	(3.4)	(1.9)	(0.6)	(0.8)	(0.9)	(1.2)	(1.2)	(100.0)				
190～	75	101	109	127	124	119	98	107	87	240	2	129	88	48	39	8	10	25	32	26	1,594	21	1,615	193.55	187.54
(%)	(4.7)	(6.3)	(6.8)	(8.0)	(7.8)	(7.5)	(6.1)	(6.7)	(5.5)	(15.1)	(0.1)	(8.1)	(5.5)	(3.0)	(2.4)	(0.5)	(0.6)	(1.6)	(2.0)	(1.6)	(100.0)				
210～	57	77	63	84	63	70	77	55	58	124	2	93	62	37	20	5	13	15	14	16	1,005	11	1,016	196.82	229.24
(%)	(5.7)	(7.7)	(6.3)	(8.4)	(6.3)	(7.0)	(7.7)	(5.5)	(5.8)	(12.3)	(0.2)	(9.3)	(6.2)	(3.7)	(2.0)	(0.5)	(1.3)	(1.5)	(1.4)	(1.6)	(100.0)				
合計	7,270	10,503	11,313	12,015	12,244	12,121	11,527	10,572	9,676	21,613	274	12,657	7,253	4,188	2,415	585	974	1,051	1,644	1,285	151,180	1,699	152,879	171.60	161.90
(%)	(4.8)	(6.9)	(7.5)	(7.9)	(8.1)	(8.0)	(7.6)	(7.0)	(6.4)	(14.3)	(0.2)	(8.4)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
記載なし	4,603	6,443	6,703	7,215	7,369	7,037	6,695	6,235	5,497	12,900	170	7,182	4,273	2,496	1,510	379	590	663	1,155	877	89,992	4,685	94,677	173.13	167.32
(%)	(5.1)	(7.2)	(7.4)	(8.0)	(8.2)	(7.8)	(7.4)	(6.9)	(6.1)	(14.3)	(0.2)	(8.0)	(4.7)	(2.8)	(1.7)	(0.4)	(0.7)	(0.7)	(1.3)	(1.0)	(100.0)				
総計	11,873	16,946	18,016	19,230	19,613	19,158	18,222	16,807	15,173	34,513	444	19,839	11,526	6,684	3,925	964	1,564	1,714	2,799	2,162	241,172	6,384	247,556	172.17	163.94
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
平均	106.23	105.68	105.32	105.71	105.21	105.80	106.05	106.43	106.31	107.19	110.36	108.46	110.20	110.98	112.09	113.95	111.56	113.78	113.58	115.08	106.99	106.99	106.99		
標準偏差	34.47	33.82	33.51	33.65	33.22	33.22	33.29	33.42	32.84	33.17	33.53	33.67	34.25	34.35	34.56	34.58	34.44	36.41	33.84	35.58	33.63	33.80	33.64		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査