

表 3513 PTH値(pg/mL) 透析前収縮期血圧(mmHg)別（透析患者全体）PTH測定法:intact-PTH

透析前収縮期血圧	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
60未満	2	2		2	3	5	1	3	3	5		2	1	1	1						31	2	33	158.65	104.81
(%)	(6.5)	(6.5)		(6.5)	(9.7)	(16.1)	(3.2)	(9.7)	(9.7)	(16.1)		(6.5)	(3.2)	(3.2)	(3.2)					(100.0)					
60～	24	33	30	27	26	27	15	28	12	49		26	19	9	8	2	3	3	10	5	356	17	373	184.38	193.95
(%)	(6.7)	(9.3)	(8.4)	(7.6)	(7.3)	(7.6)	(4.2)	(7.9)	(3.4)	(13.8)		(7.3)	(5.3)	(2.5)	(2.2)	(0.6)	(0.8)	(0.8)	(2.8)	(1.4)	(100.0)				
80～	204	202	241	227	266	211	201	170	154	364	4	193	131	76	31	15	25	14	40	45	2,814	76	2,890	174.50	222.64
(%)	(7.2)	(7.2)	(8.6)	(8.1)	(9.5)	(7.5)	(7.1)	(6.0)	(5.5)	(12.9)	(0.1)	(6.9)	(4.7)	(2.7)	(1.1)	(0.5)	(0.9)	(0.5)	(1.4)	(1.6)	(100.0)				
100～	865	1,150	1,173	1,264	1,221	1,143	1,057	1,008	966	1,999	17	1,188	597	415	259	63	120	108	202	164	14,979	396	15,375	171.10	169.92
(%)	(5.8)	(7.7)	(7.8)	(8.4)	(8.2)	(7.6)	(7.1)	(6.7)	(6.4)	(13.3)	(0.1)	(7.9)	(4.0)	(2.8)	(1.7)	(0.4)	(0.8)	(0.7)	(1.3)	(1.1)	(100.0)				
120～	2,225	3,225	3,411	3,552	3,674	3,611	3,502	3,166	2,907	6,589	92	3,767	2,208	1,279	750	165	304	329	529	412	45,697	999	46,696	172.95	167.07
(%)	(4.9)	(7.1)	(7.5)	(7.8)	(8.0)	(7.9)	(7.7)	(6.9)	(6.4)	(14.4)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.7)	(0.7)	(1.2)	(0.9)	(100.0)				
140～	3,288	4,751	5,156	5,582	5,670	5,785	5,382	4,954	4,488	10,179	124	5,856	3,287	1,846	1,073	265	414	495	718	551	69,864	1,317	71,181	170.00	155.99
(%)	(4.7)	(6.8)	(7.4)	(8.0)	(8.1)	(8.3)	(7.7)	(7.1)	(6.4)	(14.6)	(0.2)	(8.4)	(4.7)	(2.6)	(1.5)	(0.4)	(0.6)	(0.7)	(1.0)	(0.8)	(100.0)				
160～	2,420	3,504	3,819	4,056	4,141	4,051	3,899	3,529	3,206	7,258	96	4,145	2,485	1,425	794	207	291	334	560	413	50,633	875	51,508	170.93	158.94
(%)	(4.8)	(6.9)	(7.5)	(8.0)	(8.2)	(8.0)	(7.7)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.9)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
180～	943	1,359	1,354	1,557	1,595	1,517	1,411	1,329	1,173	2,727	35	1,609	961	561	338	94	120	159	227	162	19,231	363	19,594	173.73	159.19
(%)	(4.9)	(7.1)	(7.0)	(8.1)	(8.3)	(7.9)	(7.3)	(6.9)	(6.1)	(14.2)	(0.2)	(8.4)	(5.0)	(2.9)	(1.8)	(0.5)	(0.6)	(0.8)	(1.2)	(0.8)	(100.0)				
200～	236	367	386	349	403	369	348	310	320	657	12	400	248	142	82	24	46	41	70	41	4,851	79	4,930	176.36	173.96
(%)	(4.9)	(7.6)	(8.0)	(7.2)	(8.3)	(7.6)	(7.2)	(6.4)	(6.6)	(13.5)	(0.2)	(8.2)	(5.1)	(2.9)	(1.7)	(0.5)	(0.9)	(0.8)	(1.4)	(0.8)	(100.0)				
220～	42	87	75	76	78	89	70	67	57	126	2	91	35	42	18	8	13	5	14	15	1,010	24	1,034	183.79	194.45
(%)	(4.2)	(8.6)	(7.4)	(7.5)	(7.7)	(8.8)	(6.9)	(6.6)	(5.6)	(12.5)	(0.2)	(9.0)	(3.5)	(4.2)	(1.8)	(0.8)	(1.3)	(0.5)	(1.4)	(1.5)	(100.0)				
合計	10,249	14,680	15,645	16,692	17,077	16,808	15,886	14,564	13,286	29,953	382	17,277	9,972	5,796	3,354	843	1,336	1,488	2,370	1,808	209,466	4,148	213,614	171.59	162.23
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.2)	(8.0)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.9)	(100.0)				
記載なし	1,624	2,266	2,371	2,538	2,536	2,350	2,336	2,243	1,887	4,560	62	2,562	1,554	888	571	121	228	226	429	354	31,706	2,236	33,942	176.06	174.78
(%)	(5.1)	(7.1)	(7.5)	(8.0)	(8.0)	(7.4)	(7.4)	(7.1)	(6.0)	(14.4)	(0.2)	(8.1)	(4.9)	(2.8)	(1.8)	(0.4)	(0.7)	(0.7)	(1.4)	(1.1)	(100.0)				
総計	11,873	16,946	18,016	19,230	19,613	19,158	18,222	16,807	15,173	34,513	444	19,839	11,526	6,684	3,925	964	1,564	1,714	2,799	2,162	241,172	6,384	247,556	172.17	163.94
(%)	(4.9)	(7.0)	(7.5)	(8.0)	(8.1)	(7.9)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.2)	(4.8)	(2.8)	(1.6)	(0.4)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
平均	150.26	151.07	150.95	151.16	151.32	151.43	151.36	151.32	151.16	151.38	153.07	151.68	152.07	151.89	151.45	152.98	150.66	151.94	150.88	149.49	151.29	148.38	151.24		
標準偏差	25.06	24.74	24.34	24.17	24.42	23.93	23.67	23.77	23.87	23.77	23.70	23.91	23.96	24.80	24.55	26.01	26.33	24.59	25.87	26.11	24.18	25.20	24.21		

数値下のかっこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査