

表 3519 PTH値(pg/mL) 原疾患別（施設血液透析患者 透析回数週3回）PTH測定法:intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	3,071	3,774	4,054	4,513	4,497	4,459	4,236	3,952	3,662	8,453	104	4,886	2,871	1,704	1,036	271	437	481	846	735	58,042	1,138	59,180	181.57	182.52
(%)	(5.3)	(6.5)	(7.0)	(7.8)	(7.7)	(7.7)	(7.3)	(6.8)	(6.3)	(14.6)	(0.2)	(8.4)	(4.9)	(2.9)	(1.8)	(0.5)	(0.8)	(0.8)	(1.5)	(1.3)	(100.0)				
慢性腎盂腎炎	91	134	144	120	151	140	141	137	109	240	1	133	88	44	32	8	15	18	32	30	1,808	40	1,848	182.29	184.24
(%)	(5.0)	(7.4)	(8.0)	(6.6)	(8.4)	(7.7)	(7.8)	(7.6)	(6.0)	(13.3)	(0.1)	(7.4)	(4.9)	(2.4)	(1.8)	(0.4)	(0.8)	(1.0)	(1.8)	(1.7)	(100.0)				
急速進行性糸球体腎炎	53	121	128	116	118	150	127	131	85	217	1	100	49	30	16	4	5	11	9	13	1,484	43	1,527	156.81	138.43
(%)	(3.6)	(8.2)	(8.6)	(7.8)	(8.0)	(10.1)	(8.6)	(8.8)	(5.7)	(14.6)	(0.1)	(6.7)	(3.3)	(2.0)	(1.1)	(0.3)	(0.3)	(0.7)	(0.6)	(0.9)	(100.0)				
妊娠腎/妊娠中毒症	46	63	72	64	72	59	85	53	54	118	2	73	43	30	16	7	12	13	23	13	918	10	928	195.97	220.32
(%)	(5.0)	(6.9)	(7.8)	(7.0)	(7.8)	(6.4)	(9.3)	(5.8)	(5.9)	(12.9)	(0.2)	(8.0)	(4.7)	(3.3)	(1.7)	(0.8)	(1.3)	(1.4)	(2.5)	(1.4)	(100.0)				
その他の分類不能の腎炎	47	35	60	63	75	46	58	56	43	125	1	67	35	29	21	3	5	6	13	8	796	19	815	186.07	197.50
(%)	(5.9)	(4.4)	(7.5)	(7.9)	(9.4)	(5.8)	(7.3)	(7.0)	(5.4)	(15.7)	(0.1)	(8.4)	(4.4)	(3.6)	(2.6)	(0.4)	(0.6)	(0.8)	(1.6)	(1.0)	(100.0)				
多発性囊胞腎	257	442	521	535	611	517	522	498	385	999	8	607	349	208	124	22	47	46	89	53	6,840	151	6,991	175.53	157.33
(%)	(3.8)	(6.5)	(7.6)	(7.8)	(8.9)	(7.6)	(7.6)	(7.3)	(5.6)	(14.6)	(0.1)	(8.9)	(5.1)	(3.0)	(1.8)	(0.3)	(0.7)	(0.7)	(1.3)	(0.8)	(100.0)				
腎硬化症	742	1,277	1,362	1,443	1,544	1,423	1,430	1,232	1,155	2,609	30	1,494	844	470	270	75	121	119	195	119	17,954	444	18,398	168.80	148.76
(%)	(4.1)	(7.1)	(7.6)	(8.0)	(8.6)	(7.9)	(8.0)	(6.9)	(6.4)	(14.5)	(0.2)	(8.3)	(4.7)	(2.6)	(1.5)	(0.4)	(0.7)	(0.7)	(1.1)	(0.7)	(100.0)				
悪性高血圧	61	87	92	143	114	114	118	107	100	228	1	149	85	51	25	4	12	12	26	18	1,547	41	1,588	185.95	170.71
(%)	(3.9)	(5.6)	(5.9)	(9.2)	(7.4)	(7.4)	(7.6)	(6.9)	(6.5)	(14.7)	(0.1)	(9.6)	(5.5)	(3.3)	(1.6)	(0.3)	(0.8)	(0.8)	(1.7)	(1.2)	(100.0)				
糖尿病性腎症	3,879	5,851	6,000	6,358	6,347	6,227	5,819	5,520	4,855	10,777	142	5,939	3,366	1,822	1,024	244	365	421	582	353	75,891	2,014	77,905	158.82	136.12
(%)	(5.1)	(7.7)	(7.9)	(8.4)	(8.4)	(8.2)	(7.7)	(7.3)	(6.4)	(14.2)	(0.2)	(7.8)	(4.4)	(2.4)	(1.3)	(0.3)	(0.5)	(0.6)	(0.8)	(0.5)	(100.0)				
SLE腎炎	81	101	119	105	98	99	101	81	71	164	6	108	56	30	22	6	10	7	18	11	1,294	28	1,322	168.42	180.81
(%)	(6.3)	(7.8)	(9.2)	(8.1)	(7.6)	(7.7)	(7.8)	(6.3)	(5.5)	(12.7)	(0.5)	(8.3)	(4.3)	(2.3)	(1.7)	(0.5)	(0.8)	(0.5)	(1.4)	(0.9)	(100.0)				
アミロイド腎	15	25	24	25	18	32	27	13	17	34	1	9	7	2	5		1		1		256	5	261	130.80	99.68
(%)	(5.9)	(9.8)	(9.4)	(9.8)	(7.0)	(12.5)	(10.5)	(5.1)	(6.6)	(13.3)	(0.4)	(3.5)	(2.7)	(0.8)	(2.0)		(0.4)		(0.4)		(100.0)				
痛風腎	29	52	46	51	65	65	49	35	29	108	3	66	36	22	12	5		6	11	6	696	22	718	175.78	148.20
(%)	(4.2)	(7.5)	(6.6)	(7.3)	(9.3)	(9.3)	(7.0)	(5.0)	(4.2)	(15.5)	(0.4)	(9.5)	(5.2)	(3.2)	(1.7)	(0.7)		(0.9)	(1.6)	(0.9)	(100.0)				
先天性代謝異常による腎不全	11	12	10	10	9	12	13	7	5	18	1	19	6	3	3	1	2		1	1	144	3	147	175.90	190.74
(%)	(7.6)	(8.3)	(6.9)	(6.9)	(6.3)	(8.3)	(9.0)	(4.9)	(3.5)	(12.5)	(0.7)	(13.2)	(4.2)	(2.1)	(2.1)	(0.7)	(1.4)		(0.7)	(0.7)	(100.0)				
腎・尿路結核	7	6	9	14	9	10	9	6	4	24		15	6	2	2	3	1		1		128	2	130	170.51	126.65
(%)	(5.5)	(4.7)	(7.0)	(10.9)	(7.0)	(7.8)	(7.0)	(4.7)	(3.1)	(18.8)		(11.7)	(4.7)	(1.6)	(1.6)	(2.3)	(0.8)		(0.8)		(100.0)				
腎・尿路結石	18	31	21	38	39	26	31	20	19	43		27	16	14	9	3	5	2	4	4	370	13	383	184.04	265.78
(%)	(4.9)	(8.4)	(5.7)	(10.3)	(10.5)	(7.0)	(8.4)	(5.4)	(5.1)	(11.6)		(7.3)	(4.3)	(3.8)	(2.4)	(0.8)	(1.4)	(0.5)	(1.1)	(1.1)	(100.0)				
腎・尿路腫瘍	28	42	48	49	43	44	46	45	26	76	2	42	29	11	10		4	4	7	5	561	14	575	166.52	156.49
(%)	(5.0)	(7.5)	(8.6)	(8.7)	(7.7)	(7.8)	(8.2)	(8.0)	(4.6)	(13.5)	(0.4)	(7.5)	(5.2)	(2.0)	(1.8)		(0.7)	(0.7)	(1.2)	(0.9)	(100.0)				
閉塞性尿路障害	24	33	31	27	37	23	35	25	32	61		44	19	15	5	1	4	5	11	10	442	14	456	205.94	301.74
(%)	(5.4)	(7.5)	(7.0)	(6.1)	(8.4)	(5.2)	(7.9)	(5.7)	(7.2)	(13.8)		(10.0)	(4.3)	(3.4)	(1.1)	(0.2)	(0.9)	(1.1)	(2.5)	(2.3)	(100.0)				
骨髄腫	9	18	17	9	10	11	11	11	8	18		10	6	3	2					1	144	6	150	137.96	120.07
(%)	(6.3)	(12.5)	(11.8)	(6.3)	(6.9)	(7.6)	(7.6)	(7.6)	(5.6)	(12.5)		(6.9)	(4.2)	(2.1)	(1.4)					(0.7)	(100.0)				
腎形成不全	22	17	16	23	18	20	24	24	29	47		37	23	16	10	2	1	5	10	12	356	3	359	226.65	219.20
(%)	(6.2)	(4.8)	(4.5)	(6.5)	(5.1)	(5.6)	(6.7)	(6.7)	(8.1)	(13.2)		(10.4)	(6.5)	(4.5)	(2.8)	(0.6)	(0.3)	(1.4)	(2.8)	(3.4)	(100.0)				
不明	715	1,223	1,256	1,339	1,357	1,332	1,236	1,147	1,052	2,328	27	1,308	814	448	281	59	98	105	179	176	16,480	503	16,983	173.04	172.48
(%)	(4.3)	(7.4)	(7.6)	(8.1)	(8.2)	(8.1)	(7.5)	(7.0)	(6.4)	(14.1)	(0.2)	(7.9)	(4.9)	(2.7)	(1.7)	(0.4)	(0.6)	(0.6)	(1.1)	(1.1)	(100.0)				
再導入	82	90	93	78	75	98	89	72	82	140		101	50	39	24	5	18	15	19	15	1,185	20	1,205	183.04	188.19
(%)	(6.9)	(7.6)	(7.8)	(6.6)	(6.3)	(8.3)	(7.5)	(6.1)	(6.9)	(11.8)		(8.5)	(4.2)	(3.3)	(2.0)	(0.4)	(1.5)	(1.3)	(1.6)	(1.3)	(100.0)				
その他	224	295	402	391	364	364	379	331	285	601	8	364	194	131	75	13	26	34	55	39	4,575	142	4,717	169.56	172.22
(%)	(4.9)	(6.4)	(8.8)	(8.5)	(8.0)	(8.0)	(8.3)	(7.2)	(6.2)	(13.1)	(0.2)	(8.0)	(4.2)	(2.9)	(1.6)	(0.3)	(0.6)	(0.7)	(1.2)	(0.9)	(100.0)				
合計	9,512	13,729	14,525	15,514	15,671	15,271	14,586	13,503	12,107	27,428	338	15,598	8,992	5,124	3,024	736	1,189	1,310	2,132	1,622	191,911	4,675	196,586	169.97	160.97
(%)	(5.0)	(7.2)	(7.6)	(8.1)	(8.2)	(8.0)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.1)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				
記載なし																									
(%)																									
総計	9,512	13,729	14,525	15,514	15,671	15,271	14,586	13,503	12,107	27,428	338	15,598	8,992	5,124	3,024	736	1,189	1,310	2,132	1,622	191,911	4,675	196,586	169.97	160.97
(%)	(5.0)	(7.2)	(7.6)	(8.1)	(8.2)	(8.0)	(7.6)	(7.0)	(6.3)	(14.3)	(0.2)	(8.1)	(4.7)	(2.7)	(1.6)	(0.4)	(0.6)	(0.7)	(1.1)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2014年調査